

COMBINATORIAL INVARIANTS OF ALGEBRAIC HAMILTONIAN ACTIONS

IVAN V. LOSEV

ABSTRACT. To any Hamiltonian action of a reductive algebraic group G on a smooth irreducible symplectic variety X we associate certain combinatorial invariants: Cartan space, Weyl group, weight and root lattices. For cotangent bundles these invariants essentially coincide with those arising in the theory of equivariant embeddings. Using our approach we establish some properties of the latter invariants.

CONTENTS

1. Introduction	1
2. Notation and conventions	3
3. Preliminaries	4
4. Local cross-sections	5
5. Hamiltonian morphisms	7
6. Combinatorial invariants	9
7. The case of cotangent bundles	14
8. Properties of Weyl groups, root and weight lattices of G -varieties	19
References	23

1. INTRODUCTION

In this paper we study certain invariants associated with Hamiltonian actions of reductive algebraic groups. All groups and varieties are defined over an algebraically closed field $\mathbb{K}$ of characteristic zero.

Let G be a reductive algebraic group and X a Hamiltonian G -variety, i.e., a smooth symplectic variety equipped with a Hamiltonian action of G (see Section 3 for generalities on Hamiltonian actions). To define our invariants we need to fix a certain Levi subgroup $L \subset G$ and an L -stable subvariety (a so-called *L -cross-section*) $X_L \subset X$ (see Section 4 for definitions). To the pair (L, X_L) we associate the affine subspace $\mathfrak{a}_{G,X}^{(X_L)} \subset \mathfrak{z}(\mathfrak{l})$ (the *Cartan space* of X), the subgroup $W_{G,X}^{(X_L)} \subset N_G(\mathfrak{a}_{G,X}^{(X_L)})/Z_G(\mathfrak{a}_{G,X}^{(X_L)})$ (the *Weyl group* of X), and the lattices $\mathfrak{X}_{G,X}^{(X_L)}, \Lambda_{G,X}^{(X_L)} \subset \mathfrak{a}_{G,X}^{(X_L)}$ (the *weight* and *root lattices*, respectively). In the sequel, while speaking about combinatorial invariants of X , we mean these data.

Our interest to these invariants comes from the theory of equivariant embeddings of homogeneous spaces. Namely, suppose G is connected. Fix a Borel subgroup $B \subset G$ and a maximal torus $T \subset B$. To an arbitrary irreducible G -variety X_0 one associates its combinatorial invariants: the weight lattice $\mathfrak{X}_{G,X_0}$, the Cartan space $\mathfrak{a}_{G,X_0}$, the Weyl group

Key words and phrases: reductive groups, Hamiltonian actions, cotangent bundles, Weyl groups, root lattices.

2000 Mathematics Subject Classification. 14L30, 53D20.

Supported by RFBR grant 05-01-00988.

$W_{G,X_0} \subset GL(\mathfrak{a}_{G,X_0})$ and the root lattice Λ_{G,X_0} . Let us give here a short description of these invariants, precise definitions will be given in the beginning of Section 7.

By definition, the weight lattice is nothing else but the set of all weights $\lambda \in \mathfrak{X}(T)$ such that there exists a B -semiinvariant rational function on X_0 of weight λ . The Cartan space $\mathfrak{a}_{G,X_0}$ is a subspace of $\mathfrak{t}^*$ spanned by $\mathfrak{X}_{G,X_0}$. The Weyl group describes the structure of the set of so called *central* G -invariant valuations of $\mathbb{K}(X_0)$ (see [B],[K3]). Similarly, the lattice Λ_{G,X_0} introduced in [K5] is, in a certain sense, dual to the group of *central* G -equivariant automorphisms of $\mathbb{K}(X_0)$.

There is a special class of G -varieties for that the role of the above invariants is particularly important. Namely, a normal G -variety X_0 is called *spherical* if B has an open orbit on X_0 . In this case all G -valuations and all G -automorphisms are central. The Cartan space and the Weyl group play an essential role in the classification theory of spherical varieties generalizing that of toric varieties, see, for example, [K2] for details.

Two sets of invariants are closely related. Namely, to a smooth G -variety X_0 one associates its cotangent bundle X . The latter is equipped with the natural symplectic form and the action of G . This action is Hamiltonian. For L one takes $Z_G(\mathfrak{a}_{G,X_0})$. Further, Vinberg, [V2], essentially noticed that there is a distinguished choice of the subvariety X_L . It turns out that $\mathfrak{a}_{G,X_0} = \mathfrak{a}_{G,X}^{(X_L)}$, $\mathfrak{X}_{G,X_0} = \mathfrak{X}_{G,X}^{(X_L)}$, $W_{G,X_0} = W_{G,X}^{(X_L)}$, $\Lambda_{G,X_0} = \Lambda_{G,X}^{(X_L)}$ provided X_0 is quasiaffine. The first two equalities follow easily from the construction of Sections 2,3 in [K4]. The equality of the Weyl groups is the main result of [K4], the proof is rather complex. Finally, the equality for root lattices was implicitly proved in [K5].

Now we describe the content of the paper. Section 3 contains preliminaries on Hamiltonian varieties. In Section 4 we review the theory of cross-sections of Hamiltonian actions tracing back to Guillemin and Sternberg, [GS]. An example of a cross-section is provided by a subvariety X_L mentioned in the beginning of the present section.

Section 5 is devoted to the notion of a Hamiltonian morphism between two Hamiltonian varieties. Using the notion of a cross-section we define *central* automorphisms of a Hamiltonian variety. Then we study some of their properties.

In Section 6 we define the combinatorial invariants of a Hamiltonian variety and study their basic properties, Lemmas 6.9-6.15.

In Section 7 we compare the combinatorial invariants of a G -variety X_0 with those defined for the Hamiltonian action on its cotangent bundle. First of all, we recall the definitions of the combinatorial invariants for X_0 . Then we state the main result of this section, the comparison theorem 7.8. Next we define the distinguished cross-sections Σ_M of X for certain Levi subgroups $M \subset G$ and study their properties (Proposition 7.13). Finally we prove Theorem 7.8. The proof is contained in Propositions 7.15, 7.16. None of this results is actually new. Note however that our proof of the equality of the root lattices (Proposition 7.16) is much more elementary than that given in [K5] (Theorems 4.1, 7.8). The proof in [K5] was valid for arbitrary (i.e., not necessarily quasiaffine) G -varieties but used the theory of group schemes in an essential way.

In Section 8 we prove some results about W_{G,X_0} , $\mathfrak{X}_{G,X_0}$, Λ_{G,X_0} based on the comparison theorem and results of Section 6. The most important results are Propositions 8.2, 8.8 and Theorem 8.7. Proposition 8.2 allows one to compare the Weyl groups W_{G,X_0} and $W_{M,X_0/R_u(Q)}$, where Q is a parabolic subgroup of G containing B and subject to some additional conditions, and M is the Levi subgroup of Q containing T . This result plays an important role, for example, in the proof of the Knop conjecture, see [L3]. Theorem 8.7 reduces the computation of Weyl groups and weight lattices to the case when $\mathfrak{a}_{G,X_0} = \mathfrak{t}$. Proposition 8.8

continues the reduction procedure. In the case when $\mathfrak{a}_{G,X_0} = \mathfrak{t}$ it reduces the computation of $W_{G,X_0}, \Lambda_{G,X_0}$ to the case when G is simple.

Results and constructions of this paper not connected with root lattices constitute Sections 3,4 of the preprint [L2]. They were used there to construct algorithms computing Weyl groups of G -varieties. The theory of root lattices developed here will be used in a forthcoming paper to establish an algorithm computing *weight* lattices of G -varieties.

2. NOTATION AND CONVENTIONS

If an algebraic group is denoted by a capital Latin letter, then we denote its Lie algebra by the corresponding small German letter.

$\langle \alpha, v \rangle$	the pairing between elements α, v of dual vector spaces.
$A^\times$	the group of invertible elements of an associative algebra A .
$\text{Aut}^G(X)$	the group of all G -equivariant automorphisms of a G -variety X .
$\text{def}_G(X)$	the defect of an irreducible Hamiltonian G -variety X .
(G, G) (resp., $[\mathfrak{g}, \mathfrak{g}]$)	the commutant of an algebraic group G (resp., of a Lie algebra $\mathfrak{g}$)
G°	the connected component of unit of an algebraic group G .
$G *_H Y$	the homogeneous bundle over G/H with a fiber Y .
$[g, y]$	the equivalence class of (g, y) in $G *_H Y$.
G_x	the stabilizer of a point $x \in X$ under an action $G : X$.
$m_G(X)$	the maximal dimension of an orbit for an action of an algebraic group G on a variety X .
$N_G(H)$, (resp. $N_G(\mathfrak{h}), \mathfrak{n}_{\mathfrak{g}}(\mathfrak{h})$)	the normalizer of an algebraic subgroup H in an algebraic group G (resp. of a subalgebra $\mathfrak{h} \subset \mathfrak{g}$ in an algebraic group G , of a subalgebra $\mathfrak{h} \subset \mathfrak{g}$ in a Lie algebra $\mathfrak{g}$).
$N_G(H, Y)$	the stabilizer of a subset $Y \subset X$ under the action $N_G(H) : Y$, where X is a set acted on by G .
$R_u(G)$	the unipotent radical of an algebraic group G
$v(f)$	the skew-gradient of a rational function on a symplectic variety X .
$V_\lambda^{(G)}$	$= \{v \in V gv = \lambda(g)v, \forall g \in G\}$, where G is an algebraic group, V is a G -module, $\lambda \in \mathfrak{X}(G)$.
$W(\mathfrak{g})$	the Weyl group of a reductive Lie algebra $\mathfrak{g}$.
$\mathfrak{X}(G)$	the character group of an algebraic group G .
X^G	the fixed-point set for an action of G on X .
$X//G$	the categorical quotient for an action of a reductive group on an affine variety X
X^{reg}	the subset of an algebraic variety X consisting of all smooth points.
$\#X$	the cardinality of a set X .
$\mathfrak{z}(\mathfrak{g})$	the center of a Lie algebra $\mathfrak{g}$.

$Z_G(H)$, (resp. $Z_G(\mathfrak{h}), \mathfrak{z}_{\mathfrak{g}}(\mathfrak{h})$)	the centralizer of an algebraic subgroup H in an algebraic group G (resp. of a subalgebra $\mathfrak{h} \subset \mathfrak{g}$ in an algebraic group G , of a subalgebra $\mathfrak{h} \subset \mathfrak{g}$ in a Lie algebra $\mathfrak{g}$).
$\Lambda(\mathfrak{g})$	the root lattice of a reductive Lie algebra $\mathfrak{g}$.
$\mu_{G,X}$	the moment map of a Hamiltonian G -variety X .
ξ_s	the semisimple part of an element $\xi \in \mathfrak{g}$.
$\pi_{G,X}$	the quotient morphism $X \rightarrow X//G$ for the action $G : X$.

3. PRELIMINARIES

Here we recall some basic definitions and results concerning Hamiltonian actions.

In this section X is a smooth algebraic variety equipped with a regular symplectic form ω and G is a reductive algebraic group acting on X by symplectomorphisms.

Let U be an open subset of X and f a regular function on U . The *skew-gradient* $v(f)$ of f is a regular vector field on U given by the equality

$$(3.1) \quad \omega_x(v(f), \eta) = \langle d_x f, \eta \rangle, x \in U, \eta \in T_x X.$$

For $f, g \in \mathbb{K}[U]$ one can define their Poisson bracket $\{f, g\} \in \mathbb{K}[U]$ by

$$(3.2) \quad \{f, g\} = \omega(v(f), v(g)).$$

Clearly, $\{f, g\} = L_{v(f)}g$, where L denotes the Lie derivative.

To any element $\xi \in \mathfrak{g}$ one associates the velocity vector field ξ_* . Suppose there is a linear map $\mathfrak{g} \rightarrow \mathbb{K}[X], \xi \mapsto H_\xi$, satisfying the following two conditions:

- (H1) The map $\xi \mapsto H_\xi$ is G -equivariant.
- (H2) $v(H_\xi) = \xi_*$.

Definition 3.1. The action $G : X$ equipped with a linear map $\xi \mapsto H_\xi$ satisfying (H1),(H2) is said to be *Hamiltonian* and X is called a Hamiltonian G -variety.

For a Hamiltonian action $G : X$ we define the morphism $\mu_{G,X} : X \rightarrow \mathfrak{g}^*$ by the formula

$$(3.3) \quad \langle \mu_{G,X}(x), \xi \rangle = H_\xi(x), \xi \in \mathfrak{g}, x \in X.$$

This morphism is called the *moment map* of the Hamiltonian G -variety X .

Conditions (H1),(H2) are equivalent, respectively, to

- (M1) $\mu_{G,X}$ is a G -morphism.
- (M2) $\langle d_x \mu_{G,X}(v), \xi \rangle = \omega_x(\xi_* x, v)$, for all $x \in X, v \in T_x X, \xi \in \mathfrak{g}$.

Any two maps $\mu_{G,X} : X \rightarrow \mathfrak{g}^*$ satisfying conditions (M1),(M2) differ by an element of $\mathfrak{g}^{*G}$. Moreover, $H_{[\xi, \eta]} = \{H_\xi, H_\eta\} = \omega(\xi_*, \eta_*)$ (see, for example, [V2]).

The following example plays a crucial role in this paper.

Example 3.2 (Cotangent bundles). Let Y be a smooth G -variety. Let $X = T^*Y$ be the cotangent bundle of Y . X is a symplectic algebraic variety (the symplectic form is written down, for example, in [V2]). The action of G on X is Hamiltonian. The moment map is given by $\langle \mu_{G,X}((y, \alpha)), \xi \rangle = \langle \alpha, \xi_* y \rangle$. Here $y \in Y, \alpha \in T_y^* Y, \xi \in \mathfrak{g}$.

Example 3.3 (Symplectic vector spaces). Let V be a vector space equipped with a non-degenerate skew-symmetric bilinear form ω . Then V is a symplectic variety. Let G be a reductive group acting on V by linear symplectomorphisms. Then the action $G : V$ is Hamiltonian. The moment map $\mu_{G,V}$ is given by $\langle \mu_{G,V}(v), \xi \rangle = \frac{1}{2} \omega(\xi v, v), \xi \in \mathfrak{g}, v \in V$.

One fixes a G -invariant symmetric form $(\cdot, \cdot)$ on $\mathfrak{g}$ such that the restriction of $(\cdot, \cdot)$ to $\mathfrak{t}(\mathbb{Q})$ is a scalar product. Such a form is necessarily nondegenerate so we may identify $\mathfrak{g}$ and $\mathfrak{g}^*$ with respect to it. In particular, we may consider the moment map as a morphism from X to $\mathfrak{g}$.

Remark 3.4 (the restriction to a subgroup). Let H be a reductive subgroup of G and X a Hamiltonian G -variety. Then X is a Hamiltonian H -variety with the moment map $\mu_{H,X} = p \circ \mu_{G,X}$. Here p denotes the restriction map $p : \mathfrak{g}^* \rightarrow \mathfrak{h}^*$.

Finally, we recall the definition of the defect of a Hamiltonian action. The proof of the next lemma is contained, for instance, in [V2], Chapter 2, Section 2. We put $\psi_{G,X} := \pi_{G,\mathfrak{g}} \circ \mu_{G,X}$.

Lemma 3.5. *Suppose X is irreducible. The following numbers are equal:*

- (1) $\dim \ker \omega_x|_{\mathfrak{g}_{*x}}$ for $x \in X$ in general position.
- (2) $m_G(X) - m_G(\overline{\text{im } \mu_{G,X}})$.
- (3) $\dim \overline{\text{im } \psi_{G,X}}$.

These equal numbers are called the defect of X .

We denote the defect of X by $\text{def}_G(X)$.

4. LOCAL CROSS-SECTIONS

In this subsection X is a quasiprojective Hamiltonian G -variety, ω is the symplectic form on X .

The local cross-section theorem reduces the study of a Hamiltonian G -variety in an étale neighborhood of a point $x \in X$ to the case when $\mu_{G,X}(x)_s \in \mathfrak{z}(\mathfrak{g})$.

Let L be a Levi subgroup of G and $\mathfrak{l}$ the corresponding Lie algebra. Put $\mathfrak{l}^r = \{\xi \in \mathfrak{l} | \mathfrak{z}_{\mathfrak{g}}(\xi_s) \subset \mathfrak{l}\}$. It is a complement to the set of zeroes of some element from $K[\mathfrak{l}]^{N_G(\mathfrak{l})}$, see, for example, [L1], Subsection 5.1.

Put $Y = \mu_{G,X}^{-1}(\mathfrak{l}^r)$. Clearly, Y is an $N_G(L)$ -stable locally-closed subvariety of X .

Proposition 4.1 ([K6], Theorem 5.4 and [L1], Corollary 5.3 and Propositions 5.2, 5.4). *In the preceding notation*

- (1) $T_y X = \mathfrak{l}_*^\perp y \oplus T_y Y$ is a skew-orthogonal direct sum for any $y \in Y$. In particular, Y is a smooth subvariety of X and the restriction of ω to Y is nondegenerate. Thus Y is equipped with the symplectic structure.
- (2) The action $N_G(L) : Y$ is Hamiltonian with the moment map $\mu_{G,X}|_Y$.
- (3) The natural morphism $G *_ {N_G(L)} Y \rightarrow X$ is étale. Its image is saturated.

Recall that a subset Z^0 of a G -variety Z is called *saturated* if there exist a G -invariant morphism $\varphi : Z \rightarrow Z_0$ and a subset $Z_0^0 \subset Z_0$ such that $Z^0 = \varphi^{-1}(Z_0^0)$.

Definition 4.2. An irreducible (=connected) component of $\mu_{G,X}^{-1}(\mathfrak{l}^r)$ equipped with the structure of a Hamiltonian L -variety obtained by the restriction of the Hamiltonian structure from $\mu_{G,X}^{-1}(\mathfrak{l}^r)$ is called an *L -cross-section* of X .

Under some special choice of L , Proposition 4.1 can be strengthened.

Definition 4.3. The centralizer in G° of $\mu_{G,X}(x)_s$ for $x \in X$ in general position is called the *principal centralizer* of X .

The principal centralizer is defined uniquely up to G° -conjugacy (see [L1], Subsection 5.2).

Proposition 4.4 ([L1], Proposition 5.7). *Let L be the principal centralizer of X . The natural morphism $G *_{N_G(L)} \mu_{G,X}^{-1}(\mathfrak{l}^{pr}) \rightarrow X$ is an open embedding and the group $N_G(L)$ acts transitively on the set of irreducible components of $\mu_{G,X}^{-1}(\mathfrak{l}^{pr})$.*

Till the end of the subsection L is the principal centralizer and X_L is an L -cross-section of X . In the sequel we denote the open saturated subset $G\mu_{G,X}^{-1}(\mathfrak{l}^{pr}) \subset X$ by X^{pr} .

Lemma 4.5. *The following conditions are equivalent:*

- (1) $m_G(X) = \dim G$.
- (2) $\underline{\text{def}}_G(X) = \text{rk } G$.
- (3) $\overline{\text{im}} \mu_{G,X} = \mathfrak{g}$.
- (4) L is a maximal torus of G and $m_L(X_L) = \text{def}_L(X_L) = \text{rk } G$.

Proof. The equivalence of the first three conditions is quite standard, see, for example, [L1], Proposition 3.7, Corollary 3.10. It remains to show that (4) is equivalent to (1)-(3). It follows from (3) that L is a maximal torus. On the other hand, if L is a maximal torus, then (1) $\Leftrightarrow$ (4), thanks to Proposition 4.1. $\square$

Now we discuss the behavior of the principal centralizer and a corresponding cross-section under some operations with Hamiltonian varieties.

Lemma 4.6. *Let X be a Hamiltonian G -variety, L its principal centralizer and X_L its L -cross-section.*

- (1) *Let M be a Levi subgroup in G containing L . Then there is a unique M -cross-section X_M of X containing X_L , L is the principal centralizer and X_L is an L -cross-section of X_M . Conversely, any L -cross-section of X_M is also an L -cross-section of X .*
- (2) *Let X' be a Hamiltonian G -variety and $\varphi : X \rightarrow X'$ a generically finite dominant rational mapping satisfying $\mu_{G,X'} \circ \varphi = \mu_{G,X}$. Then L is the principal centralizer of X' , the rational mapping φ is defined in X_L and there exists a unique L -cross-section X'_L of X' such that $\overline{\varphi(X_L)} = \overline{X'_L}$.*
- (3) *Suppose G is connected. Let $G^0 \subset G$ be a connected subgroup containing (G, G) . Then $L^0 := G^0 \cap L$ is the principal centralizer and X_L is an L^0 -cross-section of the Hamiltonian G^0 -variety X .*
- (4) *Suppose G is connected and X satisfies the equivalent conditions of Lemma 4.5. Let $G_1, \dots, G_k$ be all simple normal subgroups of G . Then the Hamiltonian G_i -variety X also satisfies the equivalent conditions of Lemma 4.5. Put $G^{(i)} = Z(G)^\circ \prod_{j \neq i} G_j$. There exists a unique $L_i := L \cap G_i$ -cross-section X_{L_i} of X containing $G^{(i)}X_L$. Moreover, $G^{(i)}X_L$ is an open subvariety in X_{L_i} .*

Proof. *Assertion 1.* Since $\mathfrak{l}^{pr} \subset \mathfrak{m}^{pr}$, we see that X_M exists. It is unique because $\mu_{G,X}^{-1}(\mathfrak{m}^{pr})$ is smooth. Note that $\mathfrak{z}_{\mathfrak{m}}(\mu_{G,X}(x)_s) = \mathfrak{l}$ for all $x \in X_L$. It follows that L contains the principal centralizer of X_M . By assertion 3 of Proposition 4.1, $G \text{ im } \mu_{M,X_M}$ is dense in $\overline{\text{im}} \mu_{G,X}$. Therefore the principal centralizers of X_M and X are conjugate in G . Thence L is the principal centralizer of X_M . Since $\mathfrak{l}^{pr} \subset \mathfrak{m}^{pr}$, we see that X_L is an L -cross-section of X_M . Since all L -cross-sections of X_M are $N_M(L)$ -conjugate, we obtain the remaining claim.

Assertion 2. Note that $\overline{\text{im}} \mu_{G,X} = \overline{\text{im}} \mu_{G,X'}$. Thus L is the principal centralizer of X' . Since GX_L is dense in X , φ is defined in X_L . Let X'_L be a unique L -cross-section of X containing a point from $\varphi|_{X_L}$. Since $\mu_{G,X'}^{-1}(\mathfrak{l}^{pr})$ is smooth, $\overline{\varphi(X_L)} \subset \overline{X'_L}$. To prove the equality it is enough to compare dimensions using assertion 3 of Proposition 4.1.

Assertion 3. Straightforward.

Assertion 4. We may change G by some covering and assume that $G = Z(G)^\circ \times G_1 \times \dots \times G_k$. It can be easily seen that L_i is the principal centralizer of the Hamiltonian G_i -variety X . Note that $\mu_{G_i, X}$ is a $G^{(i)}$ -invariant morphism and $X_L \subset \mu_{G_i, X}^{-1}(\mathfrak{l}_i^{pr})$. Thus $G^{(i)}X_L \subset \mu_{G_i, X}^{-1}(\mathfrak{l}_i^{pr})$. Since $\mu_{G_i, X}^{-1}(\mathfrak{l}_i^{pr})$ is smooth and $G^{(i)}, X_L$ are irreducible, there exists a unique L_i -cross-section X_{L_i} of X containing $G^{(i)}X_L$. We recall that the natural morphism $G *_L X_L \rightarrow X$ is an open embedding. Thus $G^{(i)}X_L$ is a locally-closed subvariety in X and thence in X_L . Moreover, $\dim G^{(i)}X_L = \dim G^{(i)} - \text{rk } G^{(i)} + \dim X_L = \dim X - \dim G_i + \text{rk } G_i = \dim X_{L_i}$. $\square$

The following lemma is straightforward.

Lemma 4.7. *Let X be a Hamiltonian G -variety, $M \subset G$ a Levi subgroup, X_M an M -cross-section of X . Suppose that there is an action $\mathbb{K}^\times : X$ and an integer k such that $\mu_{G, X}(tx) = t^k \mu_{G, X}(x), x \in X$. Then X_M is $\mathbb{K}^\times$ -stable and $\mu_{M, X_M}(tx) = t^k \mu_{M, X_M}(x), x \in X_M$.*

5. HAMILTONIAN MORPHISMS

Definition 5.1. Let X, Y be Hamiltonian G -varieties. An etale G -equivariant morphism $\varphi : X \rightarrow Y$ is called *Hamiltonian* if it satisfies conditions 1,2 below:

- (1) $\varphi^* \omega^Y = \omega^X$. Here ω^X, ω^Y are symplectic forms on X and Y .
- (2) $\mu_{G, Y} \circ \varphi = \mu_{G, X}$.

Example 5.2. Let X_0 be a smooth G -variety and φ its automorphism. Let φ_* denote the natural symplectomorphism of T^*X_0 lifting φ . Note that the map $\varphi \mapsto \varphi_*$ is a monomorphism. If φ is a G -equivariant, then φ_* is a Hamiltonian automorphism of X .

In Section 7 we use the the following description of automorphisms from the previous example. Note that there is the natural action $\mathbb{K}^\times : X := T^*X_0$ by the fiberwise multiplication. This action satisfies the assumptions of Lemma 4.7 for $k = 1$.

Now suppose that X_0 is quasiaffine and $\phi : X \dashrightarrow X$ is a $G \times \mathbb{K}^\times$ -equivariant rational mapping preserving the symplectic form such that $\phi^*(\mathbb{K}[X]) = \mathbb{K}[X]$. Then ϕ^* induces an automorphism of $\mathbb{K}[X_0] = \mathbb{K}[X]^{\mathbb{K}^\times}$.

Lemma 5.3. *Let X, X_0, φ be such as above. Suppose that the restriction of φ^* to $\mathbb{K}[X_0]$ is induced by some regular automorphism ψ of X_0 . Then $\varphi = \psi_*$.*

Proof. Replacing φ with $\varphi\psi_*^{-1}$ we may assume that $\psi = id$. Note that φ^* preseves the Poisson bracket on X . The $\mathbb{K}[X_0]$ -module $\mathbb{K}[X]_{-1}^{(\mathbb{K}^\times)}$ is identified with the module of vector fields on X_0 . For $v \in \mathbb{K}[X_0]_{-1}^{(\mathbb{K}^\times)}, g \in \mathbb{K}[X_0]$ we have $\{v, f\} = L_v f$. It follows that φ^* is the identity on $\mathbb{K}[X_0]_{-1}^{(\mathbb{K}^\times)}$. Since X_0 is quasiaffine, there is $f \in \mathbb{K}[X_0]$ such that $\{x \in X_0 | f(x) \neq 0\}$ is affine. It follows that $\mathbb{K}[X_0]$ and $\mathbb{K}[X_0]_{-1}^{(\mathbb{K}^\times)}$ generate $\mathbb{K}(X)$. Therefore $\varphi = id$. $\square$

Now we consider a special class of Hamiltonian automorphisms. Let L be the principal centralizer and X_L an L -cross-section of X .

Definition 5.4. A Hamiltonian automorphism φ of X is said to be *central*, if $\varphi(X_L) = X_L$ and there exists $t_\varphi \in Z(L)^\circ$ such that $\varphi(x) = t_\varphi x$ for all $x \in X_L$. Central automorphisms form a group denoted by $\mathfrak{A}_{G, X}^{(\cdot)}$.

Remark 6.6 in the next section shows that the condition of being central does not depend on the choice of the principal centralizer and a cross-section.

The following lemma partially justifies the term "central".

Lemma 5.5. *Let ψ be a Hamiltonian automorphism of X fixing X_L . Then ψ commutes with any central Hamiltonian automorphism of X .*

Proof. Let φ be a central Hamiltonian automorphism of X . Since ψ, φ are G -equivariant and GX_L is dense in X , it is enough to show that $\psi|_{X_L}\varphi|_{X_L} = \varphi|_{X_L}\psi|_{X_L}$. It remains to note that the restriction of ψ to X_L is $Z(L)^\circ$ -equivariant. $\square$

Let $A_{G,X}^{(X_L)}$ denote the quotient of $Z(L)^\circ$ by the inefficiency kernel of the action $Z(L)^\circ : X_L$. The action $N_G(L, X_L)$ on $Z(L)^\circ$ descends to the action $N_G(L, X_L) : A_{G,X}^{(X_L)}$.

The restriction of a central automorphism to X_L determines the homomorphism $\mathfrak{A}_{G,X}^{(\cdot)} \rightarrow A_{G,X}^{(X_L)}$. Since GX_L is dense in X , this homomorphism is injective. Let us denote its image by $\mathfrak{A}_{G,X}^{(X_L)}$. Since the restriction of any G -equivariant automorphism to X_L is $N_G(L, X_L)$ -equivariant, we see that $\mathfrak{A}_{G,X}^{(X_L)} \subset (A_{G,X}^{(X_L)})^{N_G(L, X_L)}$.

Now we give a more convenient description of the subgroup $\mathfrak{A}_{G,X}^{(X_L)} \subset (A_{G,X}^{(X_L)})^{N_G(L, X_L)}$. Any element l of the last group defines a G -equivariant automorphism l_* of $X^{pr} = G *_{N_G(L, X_L)} X_L$ by $l_*[g, y] = [g, ly]$.

Lemma 5.6. *Let $l \in (A_{G,X}^{(X_L)})^{N_G(L, X_L)}$. Then*

- (1) *The automorphism l_* of X^{pr} is Hamiltonian and central. So $(A_{G,X}^{(X_L)})^{N_G(L, X_L)} = \mathfrak{A}_{G,X^{pr}}^{(X_L)}$.*
- (2) *$l \in \mathfrak{A}_{G,X}^{(X_L)}$ iff l_* can be extended to a morphism $X \rightarrow X$.*

Proof. The equality $\mu_{G,X^{pr}} \circ l_* = \mu_{G,X^{pr}}$ stems from $\mu_{G,X}|_{X_L} = \mu_{L,X_L}$. To complete the proof of assertion 1 it remains to verify that l_* preserves ω . By assertion 1 of Proposition 4.1, for any $y \in X_L$ there is the skew-orthogonal direct sum decomposition $T_y X = T_y X_L \oplus \mathfrak{l}_*^\perp y$. Since X_L is l_* -stable and l_* is G -equivariant, we have $d_y(l_*)T_y X_L = T_{ly} X_L$, $d_y(l_*)\mathfrak{l}_*^\perp y = \mathfrak{l}_*^\perp(ly)$. Since $y \mapsto ly$ is a symplectomorphism of X_L , we see that $d_y(l_*)$ maps $\omega_y|_{T_y X_L}$ to $\omega_{ly}|_{T_{ly} X_L}$. On the other hand, for $\xi, \eta \in \mathfrak{l}^\perp$ the equality $\omega_y(\xi_* y, \eta_* y) = (\mu_{L,X_L}(y), [\xi, \eta])$ holds. Since $y \mapsto ly$ is a Hamiltonian automorphism of X_L , it follows that $d_y(l_*)\omega_y|_{\mathfrak{l}_*^\perp y} = \omega_{ly}|_{\mathfrak{l}_*^\perp(ly)}$.

Proceed to assertion 2. If $l \in \mathfrak{A}_{G,X}^{(X_L)}$, then there is nothing to prove. Now let $l \in (A_{G,X}^{(X_L)})^{N_G(L, X_L)}$ be such that l_* can be extended to a morphism $\varphi : X \rightarrow X$. By assertion 1, $\varphi^* \omega = \omega$. Since ω is non-degenerate, we see that φ is etale. Taking into account that φ is birational, we see that it is an open embedding. But any open embedding $X \rightarrow X$ is an isomorphism (see, for example, [I], Proposition 4). $\square$

Corollary 5.7. *If X is quasiaffine, then $\mathfrak{A}_{G,X}^{(X_L)}$ is a closed subgroup of $A_{G,X}^{(X_L)}$.*

Proof. The subgroup $\mathfrak{A}_{G,X^{pr}}^{(X_L)} = (A_{G,X}^{(X_L)})^{N_G(L, X_L)} \subset A_{G,X}^{(X_L)}$ is closed. By assertion 1 of Lemma 5.6, the corollary stems from the following lemma. $\square$

Lemma 5.8. *Let G be an algebraic group acting on an open subset X^0 of a quasiaffine variety X . Then the subgroup of G consisting of all elements, that can be extended to an automorphism of X , is closed.*

Proof. $\mathbb{K}[X^0]$ is a rational G -algebra (see [PV], Lemma 1.4) and $\mathbb{K}[X]$ is a G -submodule of $\mathbb{K}[X^0]$. The set of all elements $g \in G$ such that $g\mathbb{K}[X] \subset \mathbb{K}[X]$ is given by the condition that some matrix coefficients vanish. Thence this set is closed. Thus $\{g \in G | g\mathbb{K}[X] = \mathbb{K}[X] | g\mathbb{K}[X] \subset \mathbb{K}[X], g^{-1}\mathbb{K}[X] \subset \mathbb{K}[X]\}$ is a closed subgroup of G . So we may assume that $g\mathbb{K}[X] = \mathbb{K}[X]$ for all $g \in G$.

There is a finitely generated subalgebra $A \subset \mathbb{K}[X]$ such that the corresponding map $X \rightarrow \overline{X} := \text{Spec}(A)$ is an embedding. Since $\mathbb{K}[X]$ is rational, we may change A by a

subalgebra generated by GA and assume that A is G -stable. An element $g \in G$ defines an automorphism of X iff $g(\overline{X} \setminus X) = \overline{X} \setminus X$. But the stabilizer of a subvariety is closed. $\square$

6. COMBINATORIAL INVARIANTS

Throughout this subsection X is an irreducible quasiprojective Hamiltonian G -variety. We denote by L the principal centralizer of X .

Choose an L -cross-section X_L of X . Note that the triple $(L, X_L, N_G(L, X_L))$ is defined uniquely up to G -conjugacy. Denote by T_0 the inefficiency kernel for the action $Z(L)^\circ : X_L$ and put $\mathfrak{l}_0 := \mathfrak{t}_0 \oplus [\mathfrak{l}, \mathfrak{l}]$.

Let us recall some results from [L1], Subsection 5.2. Put $\mathfrak{a}_{G,X}^{(X_L)} = \overline{\text{im } \mu_{Z(L)^\circ, X_L}}$. The restriction of $\pi_{L,\mathfrak{l}}$ to $\mathfrak{z}(\mathfrak{l}) = \mathfrak{l}^L$ is a closed embedding, so we may consider $\mathfrak{z}(\mathfrak{l})$ as a subvariety of $\mathfrak{l}/L$. Under this identification, $\mathfrak{a}_{G,X}^{(X_L)}$ is identified with $\overline{\text{im } \psi_{L,X_L}}$. It turns out that $\mathfrak{a}_{G,X}^{(X_L)}$ is an affine subspace in $\mathfrak{z}(\mathfrak{l})$ of dimension $\text{def}_G(X)$ intersecting $\mathfrak{t}_0$ in a unique point ξ_0 . Taking ξ_0 for 0 in the affine space $\mathfrak{a}_{G,X}^{(X_L)}$, we may (and will) consider $\mathfrak{a}_{G,X}^{(X_L)}$ as a vector space.

Definition 6.1. The vector space $\mathfrak{a}_{G,X}^{(X_L)}$ is called the *Cartan space* of X associated with X_L .

The affine subspace $\mathfrak{a}_{G,X}^{(X_L)} \subset \mathfrak{z}(\mathfrak{l})$ is stable under the natural action $N_G(L, X_L) : \mathfrak{z}(\mathfrak{l})$. The point ξ_0 is $N_G(L, X_L)$ -invariant. Denote the image of $N_G(L, X_L)$ in $\text{GL}(\mathfrak{a}_{G,X}^{(X_L)})$ by $W_{G,X}^{(X_L)}$. We remark that this definition of $W_{G,X}^{(X_L)}$ differs slightly from that given in [L1]. However, if G is connected, then the both definitions coincide and $W_{G,X}^{(X_L)} \cong N_G(L, X_L)/L$.

Definition 6.2. The linear group $W_{G,X}^{(X_L)}$ is said to be the *Weyl group* of the Hamiltonian G -variety X associated with X_L .

The previous definition in the case of cotangent bundles is due to Vinberg, [V1].

Remark 6.3. The affine map $\xi \mapsto \xi - \xi_0 : \mathfrak{a}_{G,X}^{(X_L)} \rightarrow \mathfrak{z}(\mathfrak{l})$ induces an isomorphism $\mathfrak{a}_{G,X}^{(X_L)} \cong \mathfrak{z}(\mathfrak{l}) \cap \mathfrak{t}_0^\perp \cong \mathfrak{z}(\mathfrak{l})/\mathfrak{t}_0$. This isomorphism is $N_G(L, X_L)/L$ -equivariant. Note that $\mathfrak{a}_{G,X}^{(X_L)}$ is identified with the Lie algebra of the group $A_{G,X}^{(X_L)} \cong Z(L)^\circ/T_0$.

Further, note that if X possesses an action of $\mathbb{K}^\times$ satisfying the assumptions of Lemma 4.7, then $0 \in \mathfrak{a}_{G,X}^{(X_L)}$ or, in other words, $\mathfrak{a}_{G,X}^{(X_L)} = \mathfrak{z}(\mathfrak{l}) \cap \mathfrak{t}_0^\perp$.

Definition 6.4. The lattice $\mathfrak{X}(A_{G,X}^{(X_L)}) \subset (\mathfrak{z}(\mathfrak{l})/\mathfrak{t}_0)^* \cong \mathfrak{z}(\mathfrak{l})/\mathfrak{t}_0 \cong \mathfrak{a}_{G,X}^{(X_L)}$ is called the *weight lattice* of the Hamiltonian G -variety X associated with X_L and is denoted by $\mathfrak{X}_{G,X}^{(X_L)}$.

Definition 6.5. The annihilator of $\mathfrak{A}_{G,X}^{(X_L)}$ in $\mathfrak{X}(A_{G,X}^{(X_L)}) = \mathfrak{X}_{G,X}^{(X_L)}$ is called the *root lattice* of the Hamiltonian G -variety X associated with X_L and is denoted by $\Lambda_{G,X}^{(X_L)}$.

Remark 6.6. Now let L' be another principal centralizer and $X_{L'}$ an L' -cross-section of X . Then there is an element $g \in G$ such that $gLg^{-1} = L'$, $gX_L = X_{L'}$. Such an element g is defined uniquely up to the right multiplication by an element from $N_G(L, X_L)$. Then $g(\mathfrak{a}_{G,X}^{(X_L)}) = \mathfrak{a}_{G,X}^{(X_{L'})}$, $gW_{G,X}^{(X_L)}g^{-1} = W_{G,X}^{(X_{L'})}$, $g\mathfrak{X}_{G,X}^{(X_L)} = \mathfrak{X}_{G,X}^{(X_{L'})}$.

Further, let φ be a Hamiltonian morphism of X satisfying the assumptions of Definition 5.4 for L, X_L . Then φ preserves $X_{L'}$ and acts on it by $gt_\varphi g^{-1}$, where t_φ is such as in Definition 5.4. It follows that $g\Lambda_{G,X}^{(X_L)} = \Lambda_{G,X}^{(X_{L'})}$.

Let us illustrate the definitions given above with two examples.

Example 6.7. Put $G = \mathrm{SL}_2$, $X = T^*(G/T)$, where T denotes the maximal torus in G consisting of diagonal matrices. As a G -variety, $X \cong G *_T (\mathbb{K}e \oplus \mathbb{K}f)$, where $e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ and $t \in T \cong \mathbb{K}^\times$ acts on $\mathbb{K}e$ by t^2 and on $\mathbb{K}f$ by t^{-2} . The moment map is given by $[g, xe + yf] \mapsto \mathrm{Ad}(g)(xe + yf)$. It is clear that $m_G(X) = \dim G$.

$$\mu_{G,X}^{-1}(\mathfrak{t}^{pr}) = \{[\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} 0 & x \\ y & 0 \end{pmatrix}] | ad = -bc = \frac{1}{2}, ya^2 = xb^2, x \neq 0\}.$$

Therefore $\mu_{G,X}^{-1}(\mathfrak{t}^{pr})$ is irreducible and $W_{G,X}^{(X_T)} = W(\mathfrak{g})$ for a unique cross-section X_T of X . The matrix $-E \in T$ acts on X trivially. The action of $T/\{\pm E\}$ on X_T is effective whence $\mathfrak{X}_{G,X}^{(X_T)} = \Lambda(\mathfrak{g})$. The group $T^{W(\mathfrak{g})}$ consists of two elements. Therefore $\Lambda_{G,X}^{(X_T)} = \Lambda(\mathfrak{g})$ or $2\Lambda(\mathfrak{g})$. Put $n = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, $h = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$. Since $nh = -hn$, we see that the translation by h is an $N_G(T)$ -automorphism of X_T . It is checked directly that $y \mapsto hy : X_T \rightarrow X_T$ is the restriction of the automorphism $[g, v] \mapsto [gn^{-1}, nv] : X \rightarrow X$. The last automorphism is Hamiltonian, see Lemma 5.6. Thus $\Lambda_{G,X}^{(X_T)} = 2\Lambda(\mathfrak{g})$.

Example 6.8. Let $X = V_1 \oplus V_2$, where V_1, V_2 are the two-dimensional irreducible symplectic SL_2 -modules. Choose a basis e_{i1}, e_{i2} of V_i . For a symplectic form on V we take a unique skew-symmetric form ω such that V_1, V_2 are isotropic and $\omega(xe_{11} + ye_{12}, ze_{21} + te_{22}) = 2 \det \begin{pmatrix} x & z \\ y & t \end{pmatrix}$. It is easily seen that $m_G(X) = \dim G$. The moment map is given by

$$\mu_{G,X}(xe_{11} + ye_{12} + ze_{21} + te_{22}) = \begin{pmatrix} xt + yz & 2yt \\ -2zx & -xt - yz \end{pmatrix}.$$

Therefore

$$\mu_{G,X}^{-1}(\mathfrak{t}^{pr}) = \{xe_{11} + ye_{12} + ze_{21} + te_{22} | yt = zx = 0, xt + yz \neq 0\}.$$

The variety $\mu_{G,X}^{-1}(\mathfrak{t}^{pr})$ has two connected components $\{xe_{11} + ye_{12} + ze_{21} + te_{22} | y = z = 0, xt \neq 0\}$ and $\{xe_{11} + ye_{12} + ze_{21} + te_{22} | x = t = 0, yz \neq 0\}$. Take the first one for X_T . One gets $W_{G,X}^{(X_T)} = \{1\}$, $\mathfrak{X}_{G,X}^{(X_T)} = \mathfrak{X}_G$. The automorphism $(v_1, v_2) \mapsto (tv_1, t^{-1}v_2)$, $v_1 \in V_1, v_2 \in V_2, t \in \mathbb{K}^\times$ is central for any $t \in \mathbb{K}^\times$. Thence $\Lambda_{G,X}^{(X_T)} = \{0\}$.

Our next task is to establish some properties of $\mathfrak{a}_{G,X}^{(\cdot)}, W_{G,X}^{(\cdot)}, \Lambda_{G,X}^{(\cdot)}, \mathfrak{X}_{G,X}^{(\cdot)}$.

Lemma 6.9. *Let L, X_L be such as above. Then $N_{G^\circ}(L, X_L) = N_G(L, X_L) \cap G^\circ$, $N_G(L, X_L)G^\circ = G$. Thus $W_{G^\circ, X}^{(X_L)}$ is a normal subgroup of $W_{G, X}^{(X_L)}$ and there is a natural epimorphism $G/G^\circ \rightarrow W_{G, X}^{(X_L)} / W_{G^\circ, X}^{(X_L)}$.*

Proof. This follows easily from the observation that $N_{G^\circ}(L, X_L)$ permutes the components of $\mu_{G,X}^{-1}(\mathfrak{t}^{pr})$ transitively. $\square$

The following lemma compares the quadruples $(\mathfrak{a}_{G,X}^{(X_L)}, W_{G,X}^{(X_L)}, \mathfrak{X}_{G,X}^{(X_L)}, \Lambda_{G,X}^{(X_L)})$ and $(\mathfrak{a}_{M,X_M}^{(X_L)}, W_{M,X_M}^{(X_L)}, \mathfrak{X}_{M,X_M}^{(X_L)}, \Lambda_{M,X_M}^{(X_L)})$, where M is a Levi subgroup in G containing L and X_M is a unique M -cross-section of X containing X_L .

Lemma 6.10. *Suppose G is connected. Let M, X_M be such as above. Then*

$$(1) \mathfrak{a}_{M,X_M}^{(X_L)} = \mathfrak{a}_{G,X}^{(X_L)}, \mathfrak{X}_{M,X_M}^{(X_L)} = \mathfrak{X}_{G,X}^{(X_L)}, A_{M,X_M}^{(X_L)} = A_{G,X}^{(X_L)}.$$

- (2) $N_M(L, Y) = N_G(L, Y) \cap M$. Equivalently, $W_{M, X_M}^{(X_L)} = W_{G, X}^{(X_L)} \cap M/L$.
- (3) $\Lambda_{M, X_M}^{(X_L)} \subset \Lambda_{G, X}^{(X_L)}$.

Proof. The first two assertions follow directly from Definitions 6.1, 6.2.

Proceed to assertion 3. The subvariety $X_M \subset X$ is a unique component of $\mu_{G, X}^{-1}(\mathfrak{m}^{pr})$ containing X_L . Thus X_M is $\mathfrak{A}_{G, X}^{(\cdot)}$ -stable. Let $\varphi \in \mathfrak{A}_{G, X}^{(\cdot)}$. Clearly, $\varphi|_{X_M}$ is a central Hamiltonian automorphism of X_M . The map $\varphi \mapsto \varphi|_{X_M}$ defines the homomorphism $\mathfrak{A}_{G, X}^{(\cdot)} \rightarrow \mathfrak{A}_{M, X_M}^{(\cdot)}$ such that the following diagram is commutative.

$$\begin{array}{ccc}
 \mathfrak{A}_{G, X}^{(\cdot)} & \xrightarrow{\quad} & \mathfrak{A}_{M, X_M}^{(\cdot)} \\
 \downarrow \iota_{G, X}^{(X_L)} & & \downarrow \iota_{M, X_M}^{(X_L)} \\
 A_{G, X}^{(X_L)} & \cong & A_{M, X_M}^{(X_L)}
 \end{array}$$

This completes the proof. $\square$

Lemma 6.11. *Let G be connected, X, X' Hamiltonian G -varieties and $\varphi : X \dashrightarrow X'$ a dominant generically finite G -equivariant rational mapping such that $\mu_{G, X'} \circ \varphi = \mu_{G, X}$. Let X'_L be the L -cross-section of X' described in assertion 2 of Lemma 4.6. Then $\mathfrak{a}_{G, X'}^{(X'_L)} = \mathfrak{a}_{G, X}^{(X_L)}$, $W_{G, X}^{(X_L)} \subset W_{G, X'}^{(X'_L)}$. If φ is birational, then $\mathfrak{X}_{G, X}^{(X_L)} = \mathfrak{X}_{G, X'}^{(X'_L)}$, $W_{G, X}^{(X_L)} = W_{G, X'}^{(X'_L)}$.*

Proof. This stems directly from the definition of X'_L . $\square$

- Lemma 6.12.** (1) $\mathfrak{X}_{G, X}^{(X_L)} = \mathfrak{X}_{G^0, X}^{(X_L)}$.
- (2) The lattices $\mathfrak{X}_{G, X}^{(X_L)}, \Lambda_{G, X}^{(X_L)} \subset \mathfrak{a}_{G, X}^{(X_L)}$ are $W_{G, X}^{(X_L)}$ -invariant.
- (3) $w\xi - \xi \in \Lambda_{G, X}^{(X_L)}$ for all $w \in W_{G, X}^{(X_L)}, \xi \in \mathfrak{X}_{G, X}^{(X_L)}$.

Proof. The first assertion is obvious. The second one stems from Remark 6.6. Proceed to the third assertion. An element $t \in A_{G, X}^{(X_L)}$ is $N_G(L, X_L)$ -invariant iff for all $\xi \in \mathfrak{X}_{G, X}^{(X_L)}, w \in W_{G, X}^{(X_L)}$ the equality $\langle \xi, t \rangle = \langle \xi, wt \rangle$ holds, equivalently, $\langle w^{-1}\xi - \xi, t \rangle = 0$. Since $\mathfrak{A}_{G, X}^{(X_L)} \subset (A_{G, X}^{(X_L)})^{N_G(L, X_L)}$, we are done. $\square$

Lemma 6.13. *Suppose G is connected and $0 \in \overline{\text{im } \psi_{G, X}}$. Let G^0 be a connected subgroup of G containing (G, G) . Put $L^0 = L \cap G^0$. Recall (Lemma 4.6) that L^0 is the principal centralizer and X_L is an L^0 -cross-section of the G^0 -variety X . Then the following assertions hold.*

- (1) The inclusion $N_{G^0}(L^0, X_L) \hookrightarrow N_G(L, X_L)$ induces an isomorphism $W_{G, X}^{(X_L)} \cong W_{G^0, X}^{(X_L)}$.
- (2) $\mathfrak{a}_{G, X}^{(X_L)} \cap \mathfrak{g}^0 \subset \mathfrak{a}_{G^0, X}^{(X_L)}$; $\mathfrak{a}_{G, X}^{(X_L)} \cap \mathfrak{g}^0$ is a $W_{G, X}^{(X_L)}$ -stable subspace of $\mathfrak{a}_{G^0, X}^{(X_L)}, \mathfrak{a}_{G, X}^{(X_L)}$.
- (3) $W_{G, X}^{(X_L)}$ acts trivially on $\mathfrak{a}_{G^0, X}^{(X_L)} / (\mathfrak{a}_{G, X}^{(X_L)} \cap \mathfrak{g}^0), \mathfrak{a}_{G, X}^{(X_L)} / (\mathfrak{a}_{G, X}^{(X_L)} \cap \mathfrak{g}^0)$.
- (4) The orthogonal projection $\mathfrak{g} \rightarrow \mathfrak{g}^0$ induces the projection $p : \mathfrak{a}_{G, X}^{(X_L)} \rightarrow \mathfrak{a}_{G^0, X}^{(X_L)}$, that is $W_{G, X}^{(X_L)}$ -equivariant.
- (5) $\Lambda_{G, X}^{(X_L)}, \Lambda_{G^0, X}^{(X_L)} \subset \mathfrak{a}_{G, X}^{(X_L)} \cap \mathfrak{g}^0$, and $\Lambda_{G, X}^{(X_L)} = \Lambda_{G^0, X}^{(X_L)}$ (the equality of lattices in $\mathfrak{a}_{G, X}^{(X_L)} \cap \mathfrak{g}^0$).

Proof. Assertion 1 stems directly from $N_G(L, X_L) \cap G^0 = N_{G^0}(L^0, X_L)$.

To prove the remaining assertions we may replace G by a covering and assume that $G = Z \times G^0$, where Z is a central subgroup of G with the Lie algebra $\mathfrak{g}^{0\perp}$.

Let T_0, T_0^0 denote the inefficiency kernels of the actions $Z(L)^\circ, Z(L^0)^\circ : X_L$. Clearly, $\mathfrak{t}_0^0 = \mathfrak{t}_0 \cap \mathfrak{g}^0$. Note that $\mathfrak{a}_{G,X}^{(X_L)} = \mathfrak{z}(\mathfrak{l}) \cap \mathfrak{t}_0^\perp, \mathfrak{a}_{G^0,X}^{(X_L)} = \mathfrak{z}(\mathfrak{l}^0) \cap \mathfrak{t}_0^{0\perp}$, since $0 \in \overline{\text{im } \psi_{G,X}}, 0 \in \overline{\text{im } \psi_{G^0,X}}$ (see Remark 6.3). Assertion 2 stems from the equalities $\mathfrak{a}_{G,X}^{(X_L)} \cap \mathfrak{g}^0 = \mathfrak{z}(\mathfrak{l}) \cap \mathfrak{g}^0 \cap \mathfrak{t}_0^\perp, \mathfrak{a}_{G^0,X}^{(X_L)} = \mathfrak{z}(\mathfrak{l}^0) \cap \mathfrak{t}_0^{0\perp} = \mathfrak{z}(\mathfrak{l}) \cap \mathfrak{g}^0 \cap (\mathfrak{t}_0 \cap \mathfrak{g}^0)^\perp$.

By definition, $W_{G,X}^{(X_L)}$ (resp., $W_{G^0,X}^{(X_L)}$) acts on $\mathfrak{a}_{G,X}^{(X_L)}$ (resp., on $\mathfrak{a}_{G^0,X}^{(X_L)}$) as $N_G(L, X_L)/L$ (resp., as $N_{G^0}(L^0, X_L)/L^0$). But both $N_G(L, X_L), N_{G^0}(L^0, X_L)$ preserve $\mathfrak{g}^0$ and act trivially $\mathfrak{g}/\mathfrak{g}^0$.

From the definitions of the corresponding moment maps we see that $\mu_{Z(L^0)^\circ, X_L}$ is the composition of $\mu_{Z(L)^\circ, X_L}$ and the orthogonal projection $\mathfrak{g} \rightarrow \mathfrak{g}^0$. Therefore $\mathfrak{a}_{G^0,X}^{(X_L)} = p(\mathfrak{a}_{G,X}^{(X_L)})$. Since the orthogonal projection $\mathfrak{l} \rightarrow \mathfrak{l}^0$ is $N_G(L, X_L)$ -equivariant, we see that p is $W_{G,X}^{(X_L)}$ -equivariant.

Finally, proceed to assertion 5. At first, we describe the relation between $\mathfrak{A}_{G,X}^{(X_L)}$ and $\mathfrak{A}_{G^0,X}^{(X_L)}$. Namely, choose $l \in Z(L)^\circ = Z \times Z(L^0)^\circ, l = (z, l^0)$. Let us show that the following conditions are equivalent.

- (a) The translation by l in X_L is the restriction of a G -equivariant automorphism of X .
- (b) The translation by l^0 in X_L is the restriction of a G^0 -equivariant automorphism of X .

To prove (a) $\Rightarrow$ (b) note that the translation by an element of Z in X is a G -equivariant automorphism. Conversely, let (b) hold and φ be the corresponding G^0 -equivariant automorphism. The automorphism φ is Hamiltonian and central, thanks to Lemma 5.3. The group Z acts on X by G^0 -equivariant automorphisms leaving X_L stable. Therefore φ is Z -equivariant (Lemma 5.5) whence G -equivariant.

The lattice $\Lambda_{G,X}^{(X_L)} \subset \mathfrak{X}(Z(L)^\circ)$ (resp. $\Lambda_{G^0,X}^{(X_L)} \subset \mathfrak{X}(Z(L^0)^\circ)$) is the annihilator of the group of all elements $l \in Z(L)^\circ$ (resp., $l^0 \in Z(L^0)^\circ$) satisfying (a) (resp., (b)). Note that $\mathfrak{X}(Z(L^0)^\circ)$ can be identified with the annihilator of $Z \subset Z(L)^\circ$ in $\mathfrak{X}(Z(L)^\circ)$. The equivalence of (a),(b) yields $\Lambda_{G,X}^{(X_L)} = \Lambda_{G^0,X}^{(X_L)}$ (the equality of subgroups in $\mathfrak{X}(Z(L)^\circ)$). An element of ZT_0 satisfies (a). It follows that $\Lambda_{G,X}^{(X_L)}$ and $\Lambda_{G^0,X}^{(X_L)}$ considered as lattices in $\mathfrak{a}_{G,X}^{(X_L)}, \mathfrak{a}_{G^0,X}^{(X_L)}$, respectively, lie in the orthogonal complement to $\mathfrak{z} + \mathfrak{t}_0$ in $\mathfrak{z}(\mathfrak{l})$. This orthogonal complement coincides with $\mathfrak{a}_{G,X}^{(X_L)} \cap \mathfrak{g}^0$. Since $\Lambda_{G,X}^{(X_L)}, \Lambda_{G^0,X}^{(X_L)}$ coincide as subgroups in $\mathfrak{X}(Z(L)^\circ)$, they also coincide as lattices in $\mathfrak{a}_{G,X}^{(X_L)} \cap \mathfrak{g}^0$. $\square$

Till the end of the subsection G is connected and X satisfies the equivalent conditions of Lemma 4.5. We denote by T a maximal torus of G . Let X_T be a T -cross-section of X . Then $\mathfrak{a}_{G,X}^{(X_T)} = \mathfrak{t}$ and $W_{G,X}^{(X_T)} \subset W(\mathfrak{g})$. Let $G_1, \dots, G_k$ be all simple normal subgroups of G so that $G = Z(G)^\circ G_1 \dots G_k$. Put $T_i = G \cap T_i$. Denote by $X_{T_i}, i = \overline{1, k}$ the T_i -cross-sections described in assertion 4 of Lemma 4.6.

Lemma 6.14. *We preserve the notation introduced in the previous paragraph.*

- (1) *There is the inclusion $W_{G,X}^{(X_T)} \subset \prod_{i=1}^k W_{G_i,X}^{(X_{T_i})}$ of subgroups in $W(\mathfrak{g})$. The image of $W_{G,X}^{(X_T)}$ in $\text{GL}(\mathfrak{t}_i)$ coincides with $W_{G_i,X}^{(X_{T_i})}$.*
- (2) *Suppose that the linear group $W_{G,X}^{(X_L)} \in \text{GL}(\mathfrak{t})$ is generated by reflections. Then $W_{G,X}^{(X_T)} = \prod_{i=1}^k W_{G_i,X}^{(X_{T_i})}$.*

Proof. Changing G by a covering, we may assume that $G = Z(G)^\circ \times \prod_{i=1}^k G_i$. Denote by $G^{(i)}$ the product of $Z(G)^\circ$ and $G_j, j \neq i$.

Let us show that $N_G(T, X_T) \subset Z(G)^\circ \times \prod_{i=1}^k N_{G_i}(T_i, X_{T_i})$. Choose $n \in N_G(T, X_T)$. We have to show that the projection n_i of n to G_i lies in $N_{G_i}(T_i, X_{T_i})$. Since $X_T \subset X_{T_i}$, we see that $nX_{T_i} \cap X_{T_i} \neq \emptyset$. Since X_{T_i} is $G^{(i)}$ -stable, we have $n_iX_{T_i} \cap X_{T_i} \neq \emptyset$. But the variety $\mu_{G_i, X}^{-1}(\mathfrak{t}_i^{pr})$ is smooth and $n_iX_{T_i}, X_{T_i}$ are its irreducible components. So $n_i \in N_{G_i}(T_i, X_{T_i})$.

Now let us show that the projection $N_G(T, X_T) \rightarrow N_{G_i}(T_i, X_{T_i})$ is surjective or, equivalently, that for any $g \in N_{G_i}(T_i, X_{T_i})$ there exists $h \in G^{(i)}$ such that $gh \in N_G(T, X_T)$. Recall that the subset $G^{(i)}X_T$ is dense in X_{T_i} . Thus for $y_1 \in X_T, h_1 \in G^{(i)}$ in general position there exist $y_2 \in X_T, h_2 \in G^{(i)}$ such that $gh_1y_1 = h_2y_2$. Since $g \in G_i, h_2 \in G^{(i)}$, we have $gh_2 = h_2g$. Thus $ghy_1 = y_2$, where $h = h_2^{-1}h_1$. The inclusions $\mu_{G, X}(y_1), \mu_{G, X}(y_2) \in \mathfrak{t}^{pr}$ imply $gh \in N_G(\mathfrak{t})$. Thence X_T, ghX_T are irreducible components of $\mu_{G, X}^{-1}(\mathfrak{t}^{pr})$ and $X_T \cap ghX_T \neq \emptyset$. It follows that $ghX_T = X_T$. This completes the proof of assertion 1.

Proceed to assertion 2. Let $s \in W_{G, X}^{(X_L)}$ be a reflection. Since $\mathfrak{t}_i \subset \mathfrak{t}, i = \overline{1, k}$, is s -stable, there exists $j \in \{1, \dots, k\}$ such that s acts trivially on $\mathfrak{t}_i, i \neq j$. Therefore there is $g \in G_j \cap N_G(T, X_T)$ mapping to s under the canonical epimorphism $N_G(T, X_T) \twoheadrightarrow W_{G, X}^{(X_T)}$. In other words, $s \in W_{G_j, X}^{(X_{T_j})}$. Since $W_{G, X}^{(X_T)}$ is generated by reflections, we see that $W_{G, X}^{(X_T)} = \prod_{i=1}^k \Gamma_i$, where $\Gamma_i \subset W_{G_i, X}^{(X_{T_i})}$. Since the projection $W_{G, X}^{(X_T)} \rightarrow W_{G_i, X}^{(X_{T_i})}$ is surjective, we get $\Gamma_i = W_{G_i, X}^{(T_i)}$. $\square$

Lemma 6.15. *Let G be connected, X satisfy the equivalent conditions of Lemma 4.5. Suppose that*

- (a) $W_{G, X}^{(\cdot)}$ is generated by reflections.
- (b) All fibers of $\psi_{G, X}$ have the same dimension.
- (c) If $\varphi \in \mathfrak{A}_{G, X^{pr}}^{(\cdot)}$ is such that φ and φ^{-1} are defined on all divisors contained in $X \setminus X^{pr}$, then φ is a regular automorphism of X .

Then $\Lambda_{G, X}^{(X_T)} = \bigoplus_{i=1}^k \Lambda_{G_i, X}^{(X_{T_i})}$.

Proof. By Lemma 6.13, $\Lambda_{G, X}^{(X_T)} = \Lambda_{(G, G), X}^{(X_T)}$. So we may (and will) assume that G is semisimple. Replacing G with a covering, we may assume, in addition, that $G = G_1 \times G_2 \times \dots \times G_k$. For $i = 1, \dots, k$ put $G^{(i)} = \prod_{j \neq i} G_j$. The proof of the lemma is in three steps.

Step 1. Let us check that $\mathfrak{A}_{G_i, X}^{(X_{T_i})} = T_i \cap \mathfrak{A}_{G, X}^{(X_T)}$ and

$$(6.1) \quad \iota_{G, X}^{(X_T)-1}(t) = \iota_{G_i, X}^{(X_{T_i})-1}(t), t \in \mathfrak{A}_{G_i, X}^{(X_{T_i})}.$$

Let $t \in \mathfrak{A}_{G_i, X}^{(X_{T_i})}$. Put $\varphi := \iota_{G_i, X}^{(X_{T_i})-1}(t)$. Note that $t \in (T_i)^{W_{G, X}^{(X_{T_i})}}$. By Lemma 6.14, $t \in T^{W_{G, X}^{(X_T)}}$. Put $\tilde{\varphi} := \iota_{G, X^{pr}}^{(X_T)-1}(t)$. Let us check that the rational mappings $\varphi, \tilde{\varphi} : X \dashrightarrow X$ coincide. The both rational mappings are G_i -equivariant and their restriction to $G^{(i)}X_T$ coincide with the translation by t . Since $G_iG^{(i)}X_T = GX_T = X^{pr}$, we have $\varphi = \tilde{\varphi}$. By assertion 2 of Lemma 5.6, $\tilde{\varphi} \in \mathfrak{A}_{G, X}^{(\cdot)}$, in other words, $t \in \mathfrak{A}_{G, X}^{(X_T)}$.

Conversely, choose $t \in T_i \cap \mathfrak{A}_{G, X}^{(X_T)}$. Put $\varphi := \iota_{G, X}^{(X_T)-1}(t)$. Since φ is G -equivariant, we have $\varphi(gy) = g\varphi(y) = tgy$ for $g \in G_i, y \in X_T$. Hence $\varphi(x) = tx$ for any $x \in X_{T_i}$. In particular, $\varphi \in \mathfrak{A}_{G_i, X}^{(\cdot)}$ and $t \in \mathfrak{A}_{G_i, X}^{(X_{T_i})}$.

Step 2. By the previous step, $\prod_{i=1}^k \mathfrak{A}_{G_i, X}^{(X_{T_i})} \subset \mathfrak{A}_{G, X}^{(X_T)}$, equivalently, $\Lambda_{G, X}^{(X_T)} \subset \bigoplus_{i=1}^k \Lambda_{G_i, X}^{(X_{T_i})}$. The inverse inclusion will follow if we show that $t := (t_1, \dots, t_k) \in \mathfrak{A}_{G, X}^{(X_T)}$ implies $t_i \in \mathfrak{A}_{G_i, X}^{(X_{T_i})}$. Thanks to step 1, it is enough to show that $t_i \in \mathfrak{A}_{G_i, X}^{(X_T)}$. Since $t_i \in T^{W_{G, X}^{(X_T)}}$, we have $t_i \in \mathfrak{A}_{G_i, X^{pr}}^{(X_T)}$.

Put $\varphi_i = \iota_{G, X^{pr}}^{(X_T)^{-1}}(t_i)$. The elements $\varphi_i \in \mathfrak{A}_{G, X^{pr}}^{(\cdot)}$ mutually commute and

$$(6.2) \quad \prod_{i=1}^k \varphi_i = \varphi,$$

where $\varphi := \iota_{G, X}^{(X_T)^{-1}}(t)$.

Step 3. Put $X_i^{pr} = G_i X_{T_i} \subset X$. It is an open affine G -stable subvariety of X . Further,

$$(6.3) \quad X \setminus X_i^{pr} = \psi_{G_i, X}^{-1}((\mathfrak{t}_i \setminus \mathfrak{t}_i^{pr})/W(\mathfrak{g}_i)) = \psi_{G, X} \left((\mathfrak{t} \setminus (\mathfrak{t}_i^{pr} \oplus \bigoplus_{j \neq i} \mathfrak{t}_j))/W(\mathfrak{g}) \right),$$

It follows from Lemma 6.14 that $t_i \in T_i^{W_{G_i, X}^{(X_{T_i})}}$. Applying (6.1) to (G, X_i^{pr}) , we get $\varphi_i = \iota_{G_i, X}^{(X_{T_i})^{-1}}(t_i)$. Therefore $\varphi_i, \varphi_i^{-1} \in \text{Aut}^G(X_i)$. By (6.3) and the assumption that $\widehat{\psi}_{G, X}$ is equidimensional, the divisors $X \setminus X_i^{pr}, X \setminus X_j^{pr}$ do not have the same irreducible component provided $i \neq j$. So $\varphi_j^{-1}, j \neq i$, is defined at the components of $X \setminus X_i$. Since $\varphi_i = \varphi \prod_{j \neq i} \varphi_j^{-1}$ (see 6.2), φ_i satisfies (c). By Lemma 5.5, $\varphi_i \in \mathfrak{A}_{G, X}^{(\cdot)}$, in other words, $t_i \in \mathfrak{A}_{G, X}^{(X_T)}$. $\square$

Remark 6.16. Clearly, condition (c) holds whenever X is affine. In fact, (b) also holds if X is affine, see, for example, [L2], Theorem 5.1.1.

7. THE CASE OF COTANGENT BUNDLES

In this section G is a reductive group, X_0 is a smooth irreducible G -variety, $X = T^*X_0$. Our objective here is to interpret invariants of X introduced in the previous section in terms of X_0 .

Till the otherwise is indicated, we suppose G is connected. Fix a Borel subgroup $B \subset G$ and a maximal torus $T \subset B$.

Definition 7.1. The lattice $\mathfrak{X}_{G, X_0} := \{\lambda \in \mathfrak{X}(T) | \mathbb{K}(X_0)_\lambda^{(B)} \neq \{0\}\}$ is called the *weight* lattice of the G -variety X_0 . Put $\mathfrak{a}_{G, X_0} = \mathfrak{X}_{G, X_0} \otimes_{\mathbb{Z}} \mathbb{K}$. We call the subspace $\mathfrak{a}_{G, X_0} \subset \mathfrak{t}^*$ the *Cartan space* of X_0 . The dimension of $\mathfrak{a}_{G, X_0}$ is called the *rank* of X_0 and is denoted by $\text{rk}_G(X_0)$.

Next we will define the Weyl group of X_0 . To this end we need the notion of a *central G -valuation*.

Definition 7.2. By a G -valuation of X_0 we mean a discrete $\mathbb{Q}$ -valued G -invariant valuation of $\mathbb{K}(X_0)$. A G -valuation is called *central* if it vanishes on $\mathbb{K}(X_0)^B$.

A central valuation v determines the element $\varphi_v \in \mathfrak{a}_{G, X_0}(\mathbb{Q})^*$ by $\langle \varphi_v, \lambda \rangle = v(f_\lambda)$, where $\lambda \in \mathfrak{X}_{G, X_0}$, $f_\lambda \in \mathbb{K}(X_0)_\lambda^{(B)} \setminus \{0\}$. The element φ_v is well-defined because v is central.

Recall that we have fixed a $W(\mathfrak{g})$ -invariant scalar product on $\mathfrak{t}(\mathbb{Q})$. This induces the scalar product on $\mathfrak{a}_{G, X_0}(\mathbb{Q}) \cong \mathfrak{a}_{G, X_0}(\mathbb{Q})^*$.

Theorem 7.3. (1) *The map $v \mapsto \varphi_v$ is injective. Its image is a finitely generated rational convex cone in $\mathfrak{a}_{G, X_0}(\mathbb{Q})^*$ called the central valuation cone of X_0 .*

(2) *The central valuation cone is simplicial (that is, there are linearly independent vectors $\alpha_1, \dots, \alpha_s \in \mathfrak{a}_{G, X_0}(\mathbb{Q})$ such that the cone coincides with $\{x | \langle \alpha_i, x \rangle \geq 0, i = \overline{1, s}\}$). Moreover, the reflections corresponding to its faces generate a finite group. This group is called the Weyl group of X and is denoted by W_{G, X_0} .*

(3) *The lattice $\mathfrak{X}_{G, X_0} \subset \mathfrak{a}_{G, X}(\mathbb{Q})$ is W_{G, X_0} -stable.*

The proof of the first part of the theorem is relatively easy. It is obtained (in a greater generality) in [K3], Korollare 3.6, 4.2, 5.2, 6.5. The second assertion is much more complicated. It was proved by Brion, [B], in the case when X_0 is spherical. Knop, [K3], used Brion's result to prove the assertion in the general case. Later, he gave an alternative proof in [K4]. The third assertion of Theorem 7.3 follows easily from the construction of the Weyl group in [K4].

Proceed to the definition of the root lattice of X_0 . First we recall the notion of a central automorphism. The following definition was given in [K5].

Definition 7.4. A G -automorphism φ of X_0 is said to be *central* if for any λ there exists $a_{\varphi,\lambda} \in \mathbb{K}^\times$ such that $\varphi(f) = a_{\varphi,\lambda} f$ for any $f \in \mathbb{K}(X_0)_\lambda^{(B)}$. Central automorphisms of X_0 form the group being denoted by $\mathfrak{A}_G(X_0)$.

It turns out that $\mathfrak{A}_G(X_0)$ is not a birational invariant of X_0 . However, if X_0^0 is an open G -subvariety in X_0 , then X_0^0 is $\mathfrak{A}_G(X_0)$ -stable and the restriction of elements of $\mathfrak{A}_G(X_0)$ to X_0^0 induces an embedding $\mathfrak{A}_G(X_0) \rightarrow \mathfrak{A}_G(X_0^0)$ ([K5], Theorem 5.1). It turns out that there is the maximal group among $\mathfrak{A}_G(X_0^0)$ ([K5], Theorem 5.4). We denote this group by $\mathfrak{A}_{G,X_0}$. Clearly, this is a birational invariant of X_0 .

Put $A_{G,X_0} = \text{Hom}(\mathfrak{X}_{G,X_0}, \mathbb{K}^\times)$. We define the map $\iota_{G,X_0} : \mathfrak{A}_{G,X_0} \rightarrow A_{G,X_0}$ by $\lambda(\iota_{G,X_0}(\varphi)) := a_{\varphi,\lambda}$. Clearly, ι_{G,X_0} is a well-defined group homomorphism.

Proposition 7.5 ([K5], Theorem 5.5). *The map ι_{G,X_0} is injective and its image is closed.*

Definition 7.6. The sublattice $\Lambda_{G,X} \subset \mathfrak{X}_{G,X} = \mathfrak{X}(A_{G,X})$ consisting of all elements vanishing on $\iota_{G,X}(\mathfrak{A}_{G,X})$ is called the *root lattice* of X .

Remark 7.7. Recall that we have fixed the invariant nondegenerate symmetric form $(\cdot, \cdot)$ on G . So we may consider $\mathfrak{a}_{G,X_0}$ as a subspace of $\mathfrak{g}$. We also recall that it is necessary to fix a Borel subgroup and a maximal torus in G to define $\mathfrak{a}_{G,X_0}, \mathfrak{X}_{G,X_0}, W_{G,X_0}, \Lambda_{G,X_0}$.

The main result of this section is a theorem comparing the quadruples $(\mathfrak{a}_{G,X_0}, W_{G,X_0}, \mathfrak{X}_{G,X_0}, \Lambda_{G,X_0})$ and $(\mathfrak{a}_{G,X}^{(\cdot)}, W_{G,X}^{(\cdot)}, \mathfrak{X}_{G,X}^{(\cdot)}, \Lambda_{G,X}^{(\cdot)})$ in the case when X_0 is quasiaffine. To state this theorem we need some preliminary considerations.

Now let G be not necessarily connected. Till the end of the section we suppose X_0 is quasiaffine. Put $L := Z_{G^\circ}(\mathfrak{a}_{G^\circ,X_0}), P := BL$. Since X_0 is quasiaffine, we see that $\mathfrak{a}_{G^\circ,X_0}$ is spanned by dominant weights. Therefore P is a parabolic subgroup of G .

For a parabolic subgroup $Q \subset G$ containing P and an open Q -stable subvariety $X'_0 \subset X_0$ put

$$(7.1) \quad \mathcal{N}(Q, X'_0) = \{(x, \beta) \in T^*X'_0 \mid \langle \beta, R_u(\mathfrak{q})x \rangle\} = \mu_{G,X}^{-1}(\mathfrak{q}) \cap T^*X'_0.$$

It follows directly from (7.1) that $\mathcal{N}(Q, X'_0) \subset X'$ is $N_G(Q)$ -stable.

Theorem 7.8 (The comparison theorem). *Let G, X_0, L, P be such as above. Then the following assertions hold.*

- (1) L is the principal centralizer of X .
- (2) There is an open P -subvariety $X'_0 \subset X_0$ such that there exists a unique L -cross-section Σ of X with $\Sigma \cap \mathcal{N}(P, X'_0) \neq \emptyset$.
- (3) If G is connected, then $\mathfrak{a}_{G,X_0} = \mathfrak{a}_{G,X}^{(\Sigma)}, A_{G,X_0} = A_{G,X}^{(\Sigma)}, \mathfrak{X}_{G,X_0} = \mathfrak{X}_{G,X}^{(\Sigma)}, W_{G,X_0} = W_{G,X}^{(\Sigma)}, \Lambda_{G,X_0} = \Lambda_{G,X}^{(\Sigma)}$.

Corollary 7.9. *In the notation of the previous theorem, $W_{G,X}^{(\Sigma)}$ is generated by reflections.*

If $X''_0 \subset X'_0$ are open P -subvarieties of X_0 , then $\mathcal{N}(P, X''_0) \subset \mathcal{N}(P, X'_0)$. It follows that the L -cross-section Σ is uniquely determined.

Actually, only the equality of the root lattices is a new result. The other claims essentially follow from [K4].

The proof of the theorem will be given after some auxiliary considerations.

Proposition 7.10. *Suppose G is connected. Then P coincides with the intersection of the stabilizers of all B -divisors on X . There is an open P -subvariety $X'_0 \subset X_0$ and a closed L -subvariety $S \subset X'_0$ such that the morphism $P *_L S \rightarrow X'_0$, $[p, x] \mapsto px$, $p \in P$, $x \in S$, is an isomorphism. For such a subvariety S the group L_0 is the inefficiency kernel of the action $L : S$.*

Proof. This was proved in [K3], Section 2, Lemma 3.1. $\square$

Let Q be a parabolic subgroup of G containing P and M the Levi subgroup of Q containing L . Put $\widetilde{M} = N_G(Q) \cap N_G(M)$. Clearly, $\widetilde{M}^\circ = M$, $N_G(Q) = \widetilde{M} \ltimes R_u(Q)$. In the sequel we will need to reduce the study of the action $G : X_0$ to the study of some action of $\widetilde{M}$.

Definition 7.11. We say that a triple (X'_0, Z_0, π) consisting of an open $N_G(Q) = N_G(R_u(\mathfrak{q}))$ -stable subvariety $X'_0 \subset X_0$, an $\widetilde{M}$ -variety Z_0 and an $N_G(Q)$ -morphism $\pi : X'_0 \rightarrow Z_0$ (we consider Z_0 as a $N_G(Q)$ -variety assuming that $R_u(Q)$ acts trivially on Z_0) *reduces* the action $Q : X_0$ if the following conditions are satisfied:

- (a) X'_0 is smooth.
- (b) The action $R_u(Q) : X'_0$ is free.
- (c) The morphism π is smooth and any its fiber consists of a unique $R_u(Q)$ -orbit.
- (d) Z_0 is smooth and quasiffine.

The following lemma is quite standard.

Lemma 7.12. *Let Q, M be such as above. Then there exists a triple (X'_0, Z_0, π) reducing the action $Q : X_0$.*

Proof. Put $X_0^1 = \{x \in X_0 \mid \dim R_u(Q)x = \dim R_u(Q)\}$. This is an open $N_G(Q)$ -subvariety in X_0 . From $P \subset Q$ it follows that $R_u(Q) \subset R_u(P)$. By Proposition 7.10, the action $R_u(P) : X_0$ is generically free. Thus $X_0^1 \neq \emptyset$. The action $R_u(Q) : X_0^1$ is free because $R_u(Q)$ is unipotent.

Since $R_u(Q)$ is a unipotent group, $\text{Quot}(\mathbb{K}[X_0]^{R_u(Q)}) = \mathbb{K}(X_0)^{R_u(Q)}$. There is a finitely generated subalgebra $A \subset \mathbb{K}[X_0]^{R_u(Q)}$ such that $\text{Quot}(A) = \mathbb{K}(X_0)^{R_u(Q)}$. Changing A by the subalgebra generated by $\widetilde{M}A$, we may assume that A is $\widetilde{M}$ -stable. Put $\widetilde{Z}_0 := \text{Spec}(A)$. Let π denote the morphism $X_0 \rightarrow \widetilde{Z}_0$ induced by the inclusion $A \subset \mathbb{K}[X_0]$.

There is an open $N_G(Q)$ -subvariety $X_0^2 \subset X_0^1$ such that $\pi(X_0^2) \subset \widetilde{Z}_0^{reg}$ and $\pi|_{X_0^2}$ is smooth. Since $\mathbb{K}(Z_0) = \mathbb{K}(X_0)^{R_u(Q)}$, a general fiber of $\pi|_{X_0^2}$ consists of a unique $R_u(Q)$ -orbit. In other words, there is an open subvariety $\widehat{Z}_0 \subset \widetilde{Z}_0^{reg}$ such that $\pi|_{X_0^2}^{-1}(z)$ is a $R_u(Q)$ -orbit for any $z \in \widehat{Z}_0$. Changing $\widehat{Z}_0$ by $\widetilde{M}\widehat{Z}_0$, we may assume that $\widehat{Z}_0$ is an $\widetilde{M}$ -variety. We take $\pi|_{X_0^2}^{-1}(\widehat{Z}_0)$ for X'_0 . Since $\pi|_{X'_0}$ is a smooth morphism, $Z_0 := \pi(X'_0)$ is an open subvariety in $\widehat{Z}_0$. $\square$

Proposition 7.13. *Let (X'_0, π, Z_0) be a triple reducing the action $Q : X_0$. Then*

- (1) *The natural projection $\mathcal{N}(Q, X'_0) \rightarrow X'_0$ is a vector bundle coinciding with the inverse image of the bundle $Z := T^*Z_0 \rightarrow Z_0$ under the morphism π . This observation provides the natural morphism $\widetilde{\pi} : \mathcal{N}(Q, X'_0) \rightarrow Z$.*

- (2) There is a unique M -cross-section Σ_M of the Hamiltonian G -variety X such that $\Sigma_M \cap \mathcal{N}(Q, X'_0) \neq \emptyset$. The intersection $\Sigma_M \cap \mathcal{N}(Q, X'_0)$ is dense in Σ_M and $R_u(Q)(\Sigma_M \cap \mathcal{N}(Q, X'_0))$ is dense in $\mathcal{N}(Q, X'_0)$. The subvariety $\Sigma_M \subset X$ is $\widetilde{M}$ -stable.
- (3) The restriction of $\widetilde{\pi} : \mathcal{N}(Q, X'_0) \rightarrow Z$ to $\Sigma_M \cap \mathcal{N}(Q, X'_0)$ is dominant, injective, $\widetilde{M}$ -equivariant. Moreover,

$$(7.2) \quad \mu_{M,Z} \circ \widetilde{\pi} = \mu_{M,\Sigma_M}.$$

Proof. Assertion 1 follows easily from assumption (b) of Definition 7.11.

Proceed to assertion 2. Since $\mathfrak{p} \subset \mathfrak{q}$, we see that $N(Q, X'_0) \cap \mathcal{N}(P, X''_0)$ is dense in $\mathcal{N}(P, X''_0)$ for any subset $X''_0 \subset X_0$ that can be included into a triple reducing the action $P : X_0$. From the proof of Theorem 3.2 from [K4] it follows that there is $y \in \mathcal{N}(Q, X'_0) \cap N(P, X''_0)$ such that $\mu_{G,X}(y) \in \mathfrak{a}_{G,X} \cap \mathfrak{l}^{pr}$. In particular, $\mathcal{N}(Q, X'_0) \cap \mu_{G,X}^{-1}(\mathfrak{m}^{pr}) \neq \emptyset$. It is well known that the morphism $R_u(Q) \times \mathfrak{m}^{pr} \rightarrow \mathfrak{m}^{pr} + R_u(\mathfrak{q}), (q, \xi) \mapsto q\xi$ is an isomorphism. Thus the morphism $R_u(Q) \times (\mathcal{N}(Q, X'_0) \cap \mu_{G,X}^{-1}(\mathfrak{m}^{pr})) \rightarrow \mathcal{N}(Q, X'_0), (g, x) \mapsto gx$, is an open embedding with the image $X' \cap \mu_{G,X}^{-1}(\mathfrak{m}^{pr} + R_u(\mathfrak{q}))$. It follows that $\mathcal{N}(Q, X'_0) \cap \mu_{G,X}^{-1}(\mathfrak{m}^{pr})$ is an irreducible variety. Therefore there is a unique M -cross-section Σ_M of X such that $\Sigma_M \cap \mathcal{N}(Q, X'_0) \neq \emptyset$. It remains to check that Σ_M is $\widetilde{M}$ -stable. This stems from the observation that $g\Sigma_M$ is an M -cross-section of X and $\mathcal{N}(Q, X'_0) \cap g\Sigma \neq \emptyset$ whenever $g \in \widetilde{M}$.

Proceed to assertion 3. The morphism in consideration is $\widetilde{M}$ -equivariant because so is $\widetilde{\pi}$. Let us check (7.2). It is enough to show that the projection of $\mu_{G,X}(y) \in \mathfrak{q}$ to $\mathfrak{m}$ coincides with $\mu_{M,Z}(\widetilde{\pi}(y))$ for any $y \in \mathcal{N}(Q, X'_0)$. Put $y = (x, \beta)$, where $x \in X'_0, \beta \in T_x^*X_0$. The annihilator of $R_u(\mathfrak{q})_*x$ of $T_x^*X_0$ is naturally identified with $T_{\pi(x)}^*Z_0$. By the definition of $\widetilde{\pi}$, $\widetilde{\pi}(y) = (\pi(x), \beta)$. Let ξ denote the projection of $\mu_{G,X}(y) \in \mathfrak{q}$ to $\mathfrak{m}$. Then $(\xi, \eta) = \langle \eta_*x, \beta \rangle = \langle \eta_*\pi(x), \beta \rangle = (\mu_{M,Z}(\widetilde{\pi}(y)), \eta)$ for all $\eta \in \mathfrak{m}$. (7.2) is proved.

Let us check that $\widetilde{\pi}|_{\Sigma_M \cap \mathcal{N}(Q, X'_0)}$ is injective. Recall that the action $R_u(Q) : X'_0$ is free and that $\widetilde{\pi} : \mathcal{N}(Q, X'_0) \rightarrow Z$ is fiberwise linear and induces an isomorphism of fibers. Since any fiber of $\pi = \widetilde{\pi}|_{X'_0}$ consists of one $R_u(Q)$ -orbit, a fiber of $\widetilde{\pi} : \mathcal{N}(Q, X'_0) \rightarrow Z$ has the same property. Choose $y_1, y_2 \in \Sigma_M \cap \mathcal{N}(Q, X'_0)$ such that $\widetilde{\pi}(y_1) = \widetilde{\pi}(y_2)$. It follows from the last equality that $\mu_{G,X}(y_1) = \mu_{G,X}(y_2)$ and that there exists $g \in R_u(Q)$ such that $gy_1 = y_2$. In particular, $\text{Ad}(g)\mu_{G,X}(y_1) = \mu_{G,X}(y_2)$. The latter is impossible, since $\mu_{G,X}(y_i) \in \mathfrak{m}^{pr}, i = 1, 2$.

To prove that $\widetilde{\pi}|_{\Sigma_M \cap \mathcal{N}(Q, X'_0)}$ is dominant it is enough to show that $\dim \Sigma_M = \dim Z$. But $\dim Z = 2 \dim Z_0 = 2(\dim X'_0 - \dim R_u(Q)) = \dim X - \dim G + \dim M = \dim \Sigma_M$. $\square$

Remark 7.14. The cross-section Σ_M from the previous proposition does not depend on the choice of X'_0 . Indeed, if X'_{01}, X'_{02} are subsets of X_0 included into triples reducing the action $Q : X$, then their intersection, being an open $N_G(Q)$ -subvariety of X'_{01} , possesses the same property. Now let Σ_M be the M -cross-section of X constructed by $X'_{01} \cap X'_{02}$. Then Σ_M intersects both $\mathcal{N}(Q, X'_{01})$ and $\mathcal{N}(Q, X'_{02})$ nontrivially. In the sequel the M -cross-section Σ_M will be called *distinguished*. We write Σ instead of Σ_L .

One may regard $\mathfrak{X}_{G,X}$ as a subgroup in $\mathfrak{X}(L)$. Put $L_0 = \bigcap_{\xi \in \mathfrak{X}_{G,X}} \ker \chi$.

The following proposition proves all assertions of Theorem 7.8 but the equality of the root lattices.

Proposition 7.15. L is the principal centralizer of the Hamiltonian G -variety X and L_0 is the inefficiency kernel for the action $L : \Sigma$. If G is connected, then $\mathfrak{a}_{G,X_0} = \mathfrak{a}_{G,X}^{(\Sigma)}, \mathfrak{X}_{G,X_0} = \mathfrak{X}_{G,X}^{(\Sigma)}, W_{G,X_0} = W_{G,X}^{(\Sigma)}$. Finally, Σ is an irreducible component of $\mu_{G,X}(\mathfrak{a}^{pr})$, where $\mathfrak{a}^{pr} := \mathfrak{a}_{G,X} \cap \mathfrak{l}^{pr}$.

Proof. We may assume that G is connected. Let X'_0, S satisfy the assumptions of Proposition 7.10 and $\pi : X'_0 \cong P *_L S \cong R_u(P) \times S \twoheadrightarrow S$ be the quotient morphism for the action $R_u(P) : X'_0$. Clearly, the triple (X'_0, π, S) reduces the action $P : X_0$. From Proposition 7.10 it follows that L_0 is the inefficiency kernel for the action $L : S$ and hence also for the action $L : T^*S$. By assertion 3 of Proposition 7.13, there is a birational L -equivariant rational mapping $\Sigma_L \dashrightarrow T^*S$. Therefore L_0 is the inefficiency kernel for the action $L : \Sigma_L$. Hence L is the principal centralizer for the Hamiltonian L -variety Σ_L . Applying assertion 1 of Lemma 4.6, we see that L is the principal centralizer of X . It follows from Remark 6.3 that $\mathfrak{a}_{G,X}^{(\Sigma)} = \mathfrak{l} \cap \mathfrak{l}_0^\perp = \mathfrak{a}_{G,X_0}, \mathfrak{X}_{G,X_0} = \mathfrak{X}_{G,X}^{(\Sigma)}$. The equality of the Weyl groups was essentially proved in [T], Remark 5. The last assertion is now clear. $\square$

The equality $\Lambda_{G,X_0} = \Lambda_{G,X}^{(\Sigma)}$ will follow if we prove

Proposition 7.16. *The homomorphism $\varphi \mapsto \varphi_*$ (see Lemma 5.3) induces an isomorphism $\mathfrak{A}_{G,X_0} \cong \mathfrak{A}_{G,X}^{(\cdot)}$. The following diagram is commutative:*

$$\begin{array}{ccc} \mathfrak{A}_{G,X_0} & \xrightarrow{\varphi \mapsto \varphi_*} & \mathfrak{A}_{G,X}^{(\cdot)} \\ \downarrow \iota_{G,X_0} & & \downarrow \iota_{G,X}^{(\Sigma)} \\ A_{G,X_0} & \cong & A_{G,X}^{(\Sigma)} \end{array}$$

Moreover, X satisfies condition (c) of Lemma 6.15.

The proof will be given after three auxiliary lemmas.

Lemma 7.17. *The natural embedding $\mathfrak{A}_G(X_0) \hookrightarrow \mathfrak{A}_{G,X_0}$ is an isomorphism.*

Proof. Choose $\gamma \in \mathfrak{A}_{G,X_0}$. For any rational G -algebra A the subset $A^{(B)} \subset \mathbb{K}(X_0)$ is γ -stable. Thus A is γ -stable. Choose an open G -embedding $X_0 \hookrightarrow \overline{X}_0$, where $\overline{X}_0$ is an affine G -variety, existing by Theorem 1.6 from [PV]. The subalgebra $\mathbb{K}[\overline{X}_0] \subset \mathbb{K}(X_0)$ is γ -stable. Further, any G -stable ideal $I \subset \mathbb{K}[\overline{X}_0]$ is γ -stable, since $I = \text{Span}_{\mathbb{K}}(GI^{(B)})$. Applying this observation to the ideal of all functions vanishing on $\overline{X}_0 \setminus X_0$, we see that $\gamma \in \mathfrak{A}_G(X_0)$. $\square$

Lemma 7.18. *Let (X'_0, π, Z_0) be a triple reducing the action $P : X_0$ and $\varphi \in \text{Aut}^G(X_0)$. Suppose X'_0 is γ -stable and there exists $\varphi_0 \in \text{Aut}^L(Z_0)$ such that $\varphi_0 \circ \pi = \pi \circ \varphi$. Then the following conditions are equivalent.*

- (1) $\varphi \in \mathfrak{A}_{G,X_0}$.
- (2) $\varphi_0 \in \mathfrak{A}_{L,Z_0}$.

Under conditions (1),(2), $\iota_{G,X_0}(\varphi) = \iota_{L,Z_0}(\varphi_0)$, and $\varphi_0(z) = \iota_{G,X}(\varphi)z$ for any $z \in Z_0$.

Proof. Note that π induces the T -equivariant group homomorphism $\pi^* : \mathbb{K}(X_0)^{(B)} \rightarrow \mathbb{K}(Z_0)^{(L \cap B)}$. It follows that (1) $\Leftrightarrow$ (2) and $\iota_{G,X_0}(\varphi) = \iota_{L,Z_0}(\varphi_0)$. The L -varieties Z_0 and S (see Proposition 7.10) are birationally equivalent. Therefore L_0 is the inefficiency kernel for the action $L : Z_0$ whence $A_{G,X_0} \cong A_{L,Z_0} \cong L/L_0$. From the definition of $\iota_{\bullet,\bullet}$ it follows that the automorphism $(\varphi_0 \iota_{L,Z_0}(\varphi_0)^{-1})^*$ of $\mathbb{K}(Z_0)$ acts trivially on $\mathbb{K}(Z_0)^{(L/L_0)}$. Since the field $\mathbb{K}(Z_0)$ is generated by $\mathbb{K}(Z_0)^{(L/L_0)}$, we get $\varphi_0 = \iota_{L,Z_0}(\varphi_0)$. $\square$

Lemma 7.19. *Let T_0 be a quasitorus (i.e. the direct product of a torus and a commutative finite group) acting on X_0 by G -equivariant automorphisms. Then the following assertion hold:*

- (1) *The structure of a Hamiltonian G -variety on X is the restriction of the structure of a Hamiltonian $G \times T_0$ -variety.*
- (2) *A triple (X'_0, π, Z_0) reducing the action $P \times T_0^\circ : X$ also reduces the action $P : X_0$, and $\mathcal{N}(P, X'_0) = \mathcal{N}(P \times T_0, X'_0)$.*
- (3) *The group $L \times T_0^\circ$ is the principal centralizer and Σ is the distinguished $L \times T_0^\circ$ -cross-section of the Hamiltonian $G \times T_0$ -variety X .*
- (4) *Σ is T_0 -stable and the rational mapping $\Sigma \dashrightarrow T^*Z_0$ from assertion 3 of Proposition 7.13 is T_0 -equivariant.*

Proof. Assertion 1 follows directly from Example 3.2. Put $G_1 = G \times T_0$, $P_1 = P \times T_0^\circ$, $L_1 = L \times T_0^\circ$, $\tilde{L}_1 = L \times T_0$. Note that $L_1 = L_{G_1^\circ, X_0}$, $P_1 = P_{G_1^\circ, X_0}$, $\tilde{L}_1 = N_{G_1}(P_1) \cap N_{G_1}(L_1)$. Assertion 2 follows now directly from Definition 7.11. In the proof of the remaining two assertions one may assume that (X'_0, π, Z_0) reduces the action $P_1 : X_0$. In this case $\mathcal{N}(P, X'_0) = \mathcal{N}(P_1, X'_0)$. Since $\mathfrak{l}_1^{pr} = \mathfrak{l}^{pr} \times \mathfrak{t}_0$, we see that $\mu_{G_1, X}^{-1}(\mathfrak{l}_1^{pr}) = \mu_{G, X}^{-1}(\mathfrak{l}^{pr})$. This proves assertion 3. Assertion 4 stems directly from assertion 3 of Proposition 7.13. $\square$

Proof of Proposition 7.16. Let us check that $\varphi_* \in \mathfrak{A}_{G, X}^{(\cdot)}$ for all $\varphi \in \mathfrak{A}_{G, X_0}$ and that $\iota_{G, X_0}(\varphi) = \iota_{G, X}^{(\Sigma)}(\varphi_*)$. This will follow if we check that $\varphi_*(x) = \iota_{G, X}(\varphi)x$ for all $x \in \Sigma$.

$\mathfrak{A}_{G, X_0}$ is a closed subgroup of A_{G, X_0} (Proposition 7.5). Thence $T_0 := \mathfrak{A}_{G, X_0}$ is a quasitorus. Let (X'_0, π, Z_0) be a triple reducing the action of the parabolic subgroup $P \times T_0^\circ \subset G \times T_0$ on X_0 . From Lemma 7.18 it follows that φ acts on Z_0 , equivalently, on $Z := T^*Z_0$ by the translation by $\iota_{G, X_0}(\varphi)$. Applying assertion 4 of Lemma 7.19, we get the required claim.

It remains to show that for any $\psi \in \mathfrak{A}_{G, X^{pr}}^{(\cdot)}$ such that ψ, ψ^{-1} are defined on all divisors contained in $X \setminus X^{pr}$ there is $\varphi \in \mathfrak{A}_{G, X_0}$ such that $\psi = \varphi_*$. Recall that there is the natural action $\mathbb{K}^\times : X$ (see the discussion preceding Lemma 5.3). The rational mapping $\psi : X \dashrightarrow X$ is $\mathbb{K}^\times$ -equivariant. Indeed, by Lemma 4.7, Σ is $\mathbb{K}^\times$ -stable. Since ψ is central, we see that its restriction to Σ is $\mathbb{K}^\times$ -equivariant. By assertion 3 of Proposition 4.1, $\overline{G\Sigma} = X$. Thus ψ itself is $\mathbb{K}^\times$ -equivariant. Note that $\psi(\mathbb{K}[X]) = \mathbb{K}[X]$. Let T_0 denote the subgroup of all $\psi \in \mathfrak{A}_{G, X^{pr}}^{(\cdot)}$ such that $\psi(\mathbb{K}[X]) = \mathbb{K}[X]$. By the proof of Corollary 5.7, T_0 is a closed subgroup of $\mathfrak{A}_{G, X^{pr}}^{(\cdot)}$. So T_0 is a quasitorus. The Lie algebra of $\mathfrak{A}_{G, X^{pr}}^{(\Sigma)} = (A_{G, X_0})^{W_{G, X_0}}$ is $\mathfrak{a}_{G, X_0}^{W_{G, X_0}}$. It follows from [K3], Satz 8.2, that the Lie algebra $\mathfrak{A}_{G, X_0}$ also coincides with $(\mathfrak{a}_{G, X_0})^{W_{G, X_0}}$. By above, any element $\psi \in T_0^\circ$ has the form φ_* for some $\varphi \in \mathfrak{A}_{G, X_0}$.

Since $\mathbb{K}[X_0] = \mathbb{K}[X]^{\mathbb{K}^\times}$, we see that $\psi(\mathbb{K}[X_0]) = \mathbb{K}[X_0]$ for any $\psi \in T_0$. So we have the rational action $T_0 : X_0$. Let $\{\gamma_1, \dots, \gamma_s\}$ be a finite subset of T_0 mapping surjectively onto T_0/T_0° . Denote by X_0^i the domain of definition of γ_i and put $X_0^0 = \bigcap_{i=1}^s X_0^i$. Since T_0° acts on X_0 by regular automorphism, we see that T_0 acts on X_0^0 by regular automorphisms. Since the action of T_0 commutes with that of G , we see that X_0^0 is G -stable. By Lemma 5.3, $\varphi = (\varphi|_{X_0^0})_*$ for any $\varphi \in T_0$. Let (X'_0, π, Z_0) be a triple reducing the action of a parabolic subgroup $P \times T_0^\circ \subset G \times T_0$ on X_0^0 . By assertion 4 of Lemma 7.19, the group T_0 acts on $Z := T^*Z_0$ or, equivalently, on Z_0 by translations by elements of L/L_0 . It follows from Lemma 7.18 that $\varphi|_{X_0}$ is central. $\square$

8. PROPERTIES OF WEYL GROUPS, ROOT AND WEIGHT LATTICES OF G -VARIETIES

In this section X_0 is a smooth quas affine G -variety, $X = T^*X_0$, L, L_0, P, Σ have the same meaning as in the previous section.

First of all, we note that Theorem 7.8 allows one to define $\mathfrak{a}_{G, X_0}, W_{G, X_0}, \mathfrak{X}_{G, X_0}, \Lambda_{G, X_0}$ even if G is disconnected. Namely, we put $\mathfrak{a}_{G, X_0} = \mathfrak{a}_{G, X}^{(\Sigma)}, W_{G, X_0} = W_{G, X}^{(\Sigma)}, \mathfrak{X}_{G, X_0} = \mathfrak{X}_{G, X}^{(\Sigma)}, \Lambda_{G, X_0} =$

$\Lambda_{G,X}^{(\Sigma)}$. Obviously, $\mathfrak{X}_{G,X_0} = \mathfrak{X}_{G^\circ,X_0}$, $\mathfrak{a}_{G,X_0} = \mathfrak{a}_{G^\circ,X_0}$. The following proposition establishes a relation between W_{G,X_0} , $W_{G^\circ,X}$.

- Proposition 8.1.** (1) $N_G(B) \cap N_G(T) \subset N_G(L, \Sigma)$. In particular, $N_G(B) \cap N_G(L) \subset N_G(\mathfrak{a}_{G,X_0})$.
 (2) Let Γ denote the image of $N_G(B) \cap N_G(L)$ in $\mathrm{GL}(\mathfrak{a}_{G,X_0})$. Then $W_{G^\circ,X_0} \subset W_{G,X_0}$ is a normal subgroup and $W_{G,X_0} = W_{G^\circ,X_0} \Gamma$.
 (3) If $\mathfrak{a}_{G,X_0} = \mathfrak{t}$, then $W_{G,X_0} = W_{G^\circ,X_0} \rtimes \Gamma$.

Proof. Note that $\mathbb{K}(X_0)^{(B)}$ is $N_G(B) \cap N_G(T)$ -stable whence $N_G(B) \cap N_G(T) \subset N_G(\mathfrak{a}_{G,X_0})$. Therefore $N_G(B) \cap N_G(T) \subset N_G(L) \cap N_G(B) \subset N_G(P) \cap N_G(L)$. By assertion 3 of Proposition 7.13, $N_G(B) \cap N_G(T) \subset N_G(L, \Sigma)$. Assertion 2 stems from Lemma 6.9. It remains to check that $W_{G^\circ,X_0} \cap \Gamma = \{1\}$ whenever $\mathfrak{a}_{G,X_0} = \mathfrak{t}$. Under the last assumption, W_{G°,X_0} is a subgroup of $W(\mathfrak{g})$. But $W(\mathfrak{g}) \cap \Gamma = \{1\}$ since the dominant Weyl chamber in $\mathfrak{t}(\mathbb{R})$ is Γ -stable. $\square$

Till the end of the subsection we suppose that G is connected. Let Q be a parabolic subgroup in G containing P and M the Levi subgroup in Q containing T . Further, let (X'_0, π, Z_0) be a triple reducing the action $R_u(Q) : X_0$. Being birational invariants, the subspace $\mathfrak{a}_{M,Z_0} \subset \mathfrak{t}$ and the group W_{M,Z_0} do not depend on the choice of Z_0 . We write $\mathfrak{a}_{M,X_0/R_u(Q)}$, $W_{M,X_0/R_u(Q)}$ instead of $\mathfrak{a}_{M,Z_0}$, W_{M,Z_0} .

Proposition 8.2. In the notation introduced above, $\mathfrak{a}_{M,X_0/R_u(Q)} = \mathfrak{a}_{G,X_0}$, $\mathfrak{X}_{M,X_0/R_u(Q)} = \mathfrak{X}_{G,X_0}$ and $W_{M,X_0/R_u(Q)} = W_{G,X_0} \cap M/T$.

Proof. The equality of the Cartan spaces and the weight lattices follows from $\mathbb{K}(X_0)^{(B)} = (\mathbb{K}(X_0)^{R_u(Q)})^{(B \cap M)}$.

The equality $\mathfrak{a}_{M,Z_0} = \mathfrak{a}_{G,X_0}$ implies that $P \cap M = P_{M,Z_0}$ whence there exists a triple (Z'_0, π_1, Y_0) reducing the action $P \cap M : Z_0$, Lemma 7.12. Put $X''_0 = \pi^{-1}(Z'_0)$. Let us check that the triple $(X''_0, \pi_1 \circ \pi, Y_0)$ reduces the action $P : X$. Clearly, $X''_0 \subset X_0$ is P -stable. Since $(R_u(Q))_x = \{1\}$ and $(R_u(P) \cap M)_{\pi(x)} = \{1\}$ for all $x \in X'_0$, we see that the action $R_u(P) : X_0$ is free. The variety Y_0 and the morphism $\pi_1 \circ \pi : X''_0 \rightarrow Y_0$ satisfy condition (c) of Definition 7.11.

Let Σ_M be the distinguished M -cross-section of X . Recall that there is the natural embedding $G *_{N_G(L, \Sigma)} \Sigma \hookrightarrow X$. Comparing the dimensions we see that $M\Sigma \cong M *_{M \cap N_G(L, \Sigma)} \Sigma$ is a dense subset of some M -cross-section of X . Since $\mathcal{N}(P, X''_0) \subset \mathcal{N}(Q, X'_0)$, we see that $M\Sigma \cap \mathcal{N}(Q, X'_0) \neq \emptyset$. Therefore $M\Sigma$ is a dense subset of Σ_M . In other words, Σ is an L -cross-section of the Hamiltonian M -variety Σ_M .

Let $\tilde{\pi} : \mathcal{N}(Q, X'_0) \rightarrow Z := T^*Z_0$ be the natural morphism defined in assertion 1 of Proposition 7.13. Recall that $\tilde{\pi}$ induces the birational M -equivariant rational mapping $\Sigma_M \dashrightarrow Z$. It is checked directly that $\tilde{\pi}(\mathcal{N}(X''_0, P)) = \mathcal{N}(Z'_0, P \cap M)$. By assertion 3 of Proposition 7.13, the subset $\tilde{\pi}(\Sigma)$ is dense in the distinguished L -cross-section Σ^Z of Z . By Lemma 6.11, $W_{M,Z}^{(\Sigma^Z)} = W_{M,\Sigma_M}^{(\Sigma)}$. Thanks to assertion 3 of Lemma 6.10, $W_{M,\Sigma_M}^{(\Sigma)} = W_{G,X}^{(\Sigma)} \cap M/L$. $\square$

Now we will give an interpretation of Lemma 6.13 for cotangent bundles.

Proposition 8.3. Let T_0 be a torus acting on X_0 by G -equivariant automorphisms. Put $\tilde{G} = G \times T_0$. Then $\mathfrak{a}_{\tilde{G},X_0} \cap \mathfrak{g} \subset \mathfrak{a}_{G,X_0}$ and $\mathfrak{a}_{G,X_0} \cap (\mathfrak{a}_{\tilde{G},X_0} \cap \mathfrak{g})^\perp \subset \mathfrak{a}_{G,X_0}^{W_{G,X_0}}$.

Proof. The assumptions of Lemma 6.13 hold (with G instead of G^0 and $G \times T_0$ instead of G). All assertions of the proposition follow directly from that lemma. $\square$

Note that the last proposition is a weaker version of results of Knop, [K3], Sätze 8.1, 8.2.

Our next objective is to reduce the computation of $\mathfrak{X}_{G,X_0}, W_{G,X_0}$ to the case when $\mathfrak{a}_{G,X_0} = \mathfrak{t}$. To this end we need the notion of the *distinguished* component of $X_0^{L_0^\circ}$.

Proposition 8.4. *We use the notation and the conventions of Proposition 7.10. Let L_1 be a normal subgroup of L_0 . Then there exists a unique irreducible component $\underline{X}_0 \subset X_0^{L_1}$ such that $\overline{U\underline{X}_0} = X_0$. This component coincides with $\overline{FS}$, where $F = R_u(P)^{L_1}$.*

Proof. Let Z be a component of $X_0^{L_1}$. If $Z \cap PS = \emptyset$, then $\overline{UZ} \subset X_0 \setminus PS$. Note that $S \subset X^{L_1}$. Consider a component Z of $X_0^{L_1}$ containing S . Then $X_0 = \overline{P *_L S} = \overline{R_u(P)S} \subset \overline{US} \subset \overline{UZ}$. Therefore it remains to prove that there exists a unique component of $X_0^{L_1}$ intersecting PS and that this component coincides with $\overline{FS}$. The L_1 -variety $PS \cong P *_L S$ is isomorphic to $R_u(P) \times S$, where L_1 acts trivially on S and by the conjugation on $R_u(P)$. It follows that $(PS)^{L_1} = FS$. The group F is irreducible because unipotent. Being an open subset of an irreducible variety, $P *_L S$ is also irreducible. It follows that S is irreducible. So FS is irreducible. $\square$

Definition 8.5. The component $\underline{X}_0 \subset X_0^{L_1}$ satisfying the assumptions of Proposition 8.4 is said to be *distinguished*.

The distinguished component for $L_1 = L_0$ was first considered by Panyushev in [Pa].

Let $\underline{X}_0$ be the distinguished component of $X_0^{L_0^\circ}$. Clearly, $\underline{X}_0$ is quasiaffine. Thanks to the following lemma, $\underline{X}_0$ is smooth.

Lemma 8.6. *Let Y be a quasiaffine variety acted on by a reductive group H and $y \in Y^H \cap Y^{\text{reg}}$. Then Y^H is smooth in y and $T_y(Y^H) = (T_y Y)^H$.*

Proof. Replacing Y with an affine H -variety containing Y as an open H -subvariety, we may assume that Y is affine. In this case one uses Corollary from [PV], Subsection 6.5. $\square$

Put $\underline{G} := N_G(L_0^\circ, \underline{X}_0)/L_0^\circ$. It is a reductive group acting on $\underline{X}_0$. Note that the Lie algebra $\underline{\mathfrak{g}}$ of $\underline{G}$ can be naturally identified with $\mathfrak{l}_0^\perp \cap \mathfrak{n}_{\mathfrak{g}}(\mathfrak{l}_0) = \mathfrak{g}^{L_0^\circ} \cap \mathfrak{z}(\mathfrak{l}_0)^\perp \subset \mathfrak{g}$. Further, note that $\mathfrak{t} \subset \mathfrak{n}_{\mathfrak{g}}(\underline{\mathfrak{g}})$. For a Borel (resp., Cartan) subalgebra in $\underline{\mathfrak{g}}$ we take $\mathfrak{b} \cap \underline{\mathfrak{g}}, \mathfrak{t} \cap \underline{\mathfrak{g}}$. We define the triple $(\underline{\mathfrak{a}}_{\underline{G}, \underline{X}_0}, \underline{\mathfrak{X}}_{\underline{G}, \underline{X}_0}, W_{\underline{G}, \underline{X}_0})$ according to this choice of Borel and Cartan subalgebras.

Theorem 8.7. *In the above notation, $\mathfrak{a}_{G,X_0} = \underline{\mathfrak{a}}_{\underline{G}, \underline{X}_0} = \mathfrak{t}$, $W_{G,X_0} = W_{\underline{G}, \underline{X}_0}$, $\mathfrak{X}_{G,X_0} = \mathfrak{X}_{\underline{G}^\circ, \underline{X}_0}$.*

Proof. Recall that there exist an open P -subvariety $X'_0 \subset X_0$ and a closed L -subvariety $S \subset X'_0$ such that $X'_0 \cong P *_L S$ (Proposition 7.10). Let Σ denote the distinguished L -cross-section of a Hamiltonian G -variety X .

Note that $\underline{\mathfrak{h}} = \mathfrak{p} \cap \underline{\mathfrak{g}}$. By Proposition 8.4, $S \subset \underline{X}_0$. Consider the morphism $\varphi : \underline{B} *_L/L_0^\circ S \rightarrow X'_0 \cap \underline{X}_0, [g, x] \mapsto gx, g \in \underline{B}, x \in S$. Recall that $X'_0 \cong R_u(P) \times S$. Thus $\underline{X}_0 \cap X'_0 = R_u(P)^{L_0^\circ} S \cong R_u(P)^{L_0^\circ} \times S$. It follows that φ is surjective. Since $\underline{B} = R_u(P)^{L_0^\circ} \rtimes (L/L_0^\circ)$, we see that φ is injective. Being a bijective morphism of smooth varieties, φ is an isomorphism. Thanks to the natural isomorphism $\mathbb{K}(S)^{(B \cap L)} \cong \mathbb{K}(X_0)^{(B)}$, we get $\mathfrak{X}_{G,X_0} = \mathfrak{X}_{L,S} = \mathfrak{X}_{\underline{G}^\circ, \underline{X}_0}, \mathfrak{a}_{G,X_0} = \mathfrak{a}_{L,S} = \mathfrak{l} \cap \mathfrak{l}_0^\perp = \mathfrak{t}$. Also we see that $\underline{X}_0 \cap X'_0$ can be included into a triple reducing the action $\underline{B} : \underline{X}_0$.

Let p denote the natural projection $X \rightarrow X_0$. For $x \in X_0^{L_0^\circ}$ the subspace $p^{-1}(x)^{L_0^\circ} \subset T_x^* X_0$ is naturally identified with $T_x^*(X_0^{L_0^\circ})$. Therefore the vector bundle $(T^* X_0)^{L_0^\circ} \rightarrow X_0^{L_0^\circ}$ coincides with the cotangent bundle of $X_0^{L_0^\circ}$. In particular, there is the natural embedding

$\underline{X} := T^* \underline{X}_0 \hookrightarrow X$. Let us check that the following diagram is commutative.

$$(8.1) \quad \begin{array}{ccc} \underline{X} & \xrightarrow{\hookrightarrow} & X \\ \downarrow \mu_{\underline{G}, \underline{X}} & & \downarrow \mu_{G, X} \\ \underline{\mathfrak{g}} & \xrightarrow{\hookrightarrow} & \mathfrak{g} \end{array}$$

Since $\mu_{G, X}$ is G -equivariant, we see that $\mu_{G, X}(\underline{X}) \subset \mathfrak{g}^{L_0^\circ}$. By the definition of $\underline{X}_0$, the group L_0° acts trivially on $\underline{X}$. Since $\underline{X}_0 \subset \underline{X} \cap \mu_{G, X}^{-1}(0)$, we see that $\mu_{G, X}(\underline{X}) \subset \mathfrak{l}_0^\perp$. It follows that $\mu_{G, X}(\underline{X}) \subset \underline{\mathfrak{g}}$. The commutativity of the diagram now stems directly from the definitions of the moment maps for the actions $\underline{G} : \underline{X}$, $G : X$ (Example 3.2).

Since $\underline{\mathfrak{h}} = \underline{\mathfrak{g}} \cap \mathfrak{p}$, (8.1) yields

$$(8.2) \quad \mathcal{N}(\underline{B}, \underline{X}_0 \cap X'_0) \subset \mathcal{N}(P, X'_0).$$

From the equality $\mathfrak{a}_{\underline{G}, \underline{X}_0} = \underline{\mathfrak{t}}$ it follows that L/L_0° is the principal centralizer of the Hamiltonian $\underline{G}$ -variety $\underline{X}$. By Proposition 7.15,

$$(8.3) \quad \Sigma \subset X^{L_0^\circ}.$$

Put $\mathfrak{a}^{pr} := \mathfrak{a}_{G, X_0} \cap \mathfrak{l}^{pr}$. By Proposition 7.15, Σ is an irreducible component of $\mu_{G, X}^{-1}(\mathfrak{a}^{pr})$ and the distinguished L/L_0° -cross-section $\underline{\Sigma}$ of $\underline{X}$ is an irreducible component of $\mu_{\underline{G}, \underline{X}}^{-1}(\mathfrak{a}^{pr})$. This observation and (8.2) yield $\underline{\Sigma} \subset \Sigma$. From $\underline{\Sigma} \subset \Sigma$, (8.3) and the smoothness of $X^{L_0^\circ}$ we deduce that $\Sigma \subset \underline{X}$. It follows from (8.1) that $\Sigma \subset \mu_{\underline{G}, \underline{X}}^{-1}(\mathfrak{a}^{pr})$. Hence $\underline{\Sigma} = \Sigma$.

Let us check the equality of the Weyl groups. We have to prove that

$$(8.4) \quad N_{\underline{G}}(L/L_0^\circ, \Sigma) = N_G(L, \Sigma)/L_0^\circ.$$

Since L_0 is the inefficiency kernel of the action $L : \Sigma$, we get $N_G(L, \Sigma) \subset N_G(L_0^\circ)$. Being a unique component of $X^{L_0^\circ}$ containing Σ , $\underline{X}$ is $N_G(L, \Sigma)$ -stable. Therefore $N_G(L, \Sigma) \subset N_G(L_0^\circ, \underline{X}) = N_G(L_0^\circ, \underline{X}_0)$. Clearly,

$$(8.5) \quad N_{\underline{G}}(L/L_0^\circ, \Sigma) = N_{N_G(L_0^\circ, \underline{X}_0)}(L, \Sigma)/L_0^\circ.$$

The inclusion $N_G(L, \Sigma) \subset N_G(L_0^\circ, \underline{X}_0)$ and (8.5) yield (8.4). $\square$

Now let $G_1, \dots, G_k$ be all simple normal subgroups in G so that $G = Z(G)^\circ G_1 \dots G_k$. We take $B_i := B \cap G_i$, $T_i := T \cap G_i$ for a Borel subgroup and a maximal torus of G_i .

Proposition 8.8. *Suppose $\mathfrak{a}_{G, X_0} = \mathfrak{t}$. Then $\mathfrak{a}_{G_i, X_0} = \mathfrak{t}_i$, $W_{G, X_0} = \prod_{i=1}^k W_{G_i, X_0}$, and $\Lambda_{G, X_0} = \bigoplus_{i=1}^k \Lambda_{G_i, X_0}$.*

Proof. Replacing G with a covering, we may assume that $G = Z(G)^\circ \times G_1 \times \dots \times G_k$. For $i = 0, k$ put $B_i = G_i \cap B$, $G^{(i)} = Z(G)^\circ \times \prod_{j \neq i} G_j$, $B^{(i)} = B \cap G^{(i)}$, $T^{(i)} = T \cap G^{(i)}$.

It follows from Proposition 7.15 and assertion 4 of Lemma 4.6 that $\mathfrak{a}_{G_i, X} = \mathfrak{t}_i$.

Thanks to Proposition 7.10, there is an open B -subvariety $X'_0 \subset X_0$ and a closed T -subvariety $S \subset X'_0$ such that $X'_0 \cong B *_T S$. For $i = 1, \dots, k$ put $Z_{0i} = B^{(i)} S$. Since $X'_0 \cong B *_T S$, we see that $Z_{0i} \cong B^{(i)} *_T S$, $X'_0 \cong B_i *_T Z_{0i}$. Therefore the subset $X'_0 \subset X_0$ can be included into a triple reducing the action $B_i : X$.

It follows immediately from (7.1) that

$$(8.6) \quad \mathcal{N}(B, X'_0) = \bigcap_{i=1}^k \mathcal{N}(B_i, X'_0).$$

Let Σ be the distinguished T -cross-section of X and Σ_i a unique T_i -cross-section of X containing $G^{(i)}\Sigma$, see assertion 4 of Lemma 4.6. Therefore the proposition will follow if we check the equality $\Sigma_i \cap \mathcal{N}(B_i, X'_0) \neq \emptyset$ and the assumptions of Lemmas 6.14, 6.15. The former inequality stems directly from $\Sigma \cap \mathcal{N}(B, X^0) \neq \emptyset$ and (8.6). By Corollary 7.9, $W_{G,X}^{(\Sigma)}$ is generated by reflections. The morphism $\psi_{G,X} : X \rightarrow \mathfrak{g}/G$ is equidimensional, thanks to Satz 6.6 from [K1]. Finally, condition (c) of Lemma 6.15 follows from Proposition 7.16. $\square$

REFERENCES

- [B] M. Brion. *Vers une généralisation des espaces symétriques*. J. Algebra, 134(1990), 115-143.
- [GS] V. Guillemin, Sh. Sternberg, *Symplectic techniques in physics*. Cambridge University Press, 1984.
- [I] S. Iitaka. *On logarithmic Kodaira dimension of algebraic varieties*. Complex analysis and algebraic geometry, p. 175-189. Iwanami Shoten, Tokyo, 1977.
- [K1] F. Knop. *Weylgruppe und Momentabbildung*. Invent. Math. 1990. V. 99. p. 1-23.
- [K2] F. Knop. *The Luna-Vust theory of spherical embeddings*. Proceedings of the Hyderabad conference on algebraic groups. Madras: Manoj Prokashan 1991.
- [K3] F. Knop. *Über Bewertungen, welche unter einer reductiven Gruppe invariant sind*. Math. Ann., 295(1993), p. 333-363.
- [K4] F. Knop. *The asymptotic behaviour of invariant collective motion*. Invent. Math. 1994. V.114. p. 309-328.
- [K5] F. Knop. *Automorphisms, root systems and compactifications*. J. Amer. Math. Soc. 9(1996), n.1, p. 153-174.
- [K6] F. Knop. *Weyl groups of Hamiltonian manifolds, I*. Preprint (1997). dg-ga/9712010.
- [L1] I.V. Losev. *Algebraic Hamiltonian actions*. Preprint (2006), arXiv:math.AG/0601023.
- [L2] I.V. Losev. *Computation of Weyl groups of G -varieties*. Preprint(2006), arXiv:math.AG/0612559.
- [L3] I.V. Losev. *Proof of the Knop conjecture*. Preprint (2006), arXiv:math.AG/0612561.
- [PV] V.L. Popov, E.B. Vinberg. *Invariant theory*. Itogi nauki i techniki. Sovr. probl. matem. Fund. napr., v. 55. Moscow, VINITI, 1989, 137-309 (in Russian). English translation in: Algebraic geometry 4, Encyclopaedia of Math. Sciences, vol. 55, Springer Verlag, Berlin, 1994.
- [Pa] D.I. Panyushev. *A restriction theorem and Poincare series for U -invariants*. Math. Annalen, 301(1995), p. 655-675.
- [T] D.A. Timashev, *Equivariant symplectic geometry of cotangent bundles II*. Moscow Math. J., v.6(2006), N2, p. 389-404.
- [V1] E.B. Vinberg. *Equivariant symplectic geometry of cotangent bundles*. Moscow Math. J. 1(2001), no. 2, 287-299.
- [V2] E.B. Vinberg.: *Commutative homogeneous spaces and coisotropic symplectic actions*. Usp. Mat. Nauk 56(2001), 3-62 (in Russian). English translation in: Russ. Math. Surveys 56(2001), 1-60.

Chair of Higher Algebra, Department of Mechanics and Mathematics, Moscow State University.

E-mail address: ivanlosev@yandex.ru